arXiv:math/0703686v1 [math.NT] 23 Mar 2007

ON UNIFORM LOWER BOUND OF THE GALOIS IMAGES ASSOCIATED TO ELLIPTIC CURVES

KEISUKE ARAI

Abstract. Let p be a prime and K be a number field. Let $\rho_{E,p} : \mathrm{G}_K \longrightarrow \mathrm{Aut}(T_pE) \cong \mathrm{GL}_2(\mathbb{Z}_p)$ be the Galois representation given by the Galois action on the p-adic Tate module of an elliptic curve E over K. Serre showed that the image of $\rho_{E,p}$ is open if E has no complex multiplication. For an elliptic curve E over K whose j-invariant does not appear in an exceptional finite set, we give an explicit uniform lower bound of the size of the image of $\rho_{E,p}$.

1. Introduction

Let k be a field of characteristic 0, and let $\mathrm{G}_k = \mathrm{Gal}(\overline{k}/k)$ be the absolute Galois group of k where $\overline{k}$ is an algebraic closure of k. Let p be a prime number. For an elliptic curve E over k, let T_pE be the p-adic Tate module of E, and let

$$\rho_{E,p} : \mathrm{G}_k \longrightarrow \mathrm{Aut}(T_pE) \cong \mathrm{GL}_2(\mathbb{Z}_p)$$

be the p-adic representation determined by the action of G_k on T_pE. By a number field we mean a finite extension of $\mathbb{Q}$.

We recall a famous theorem proved by Serre:

Theorem 1.1. *([17, IV-11]) Let K be a number field, E be an elliptic curve over K with no complex multiplication and p be a prime. Then the representation $\rho_{E,p}$ has an open image i.e. there exists a positive integer n depending on K, E and p such that*

$$\rho_{E,p}(\mathrm{G}_K) \supseteq 1 + p^n\mathrm{M}_2(\mathbb{Z}_p).$$

In this paper, we show that there exists a uniform bound of such n if we let E vary for fixed K and p.

Theorem 1.2. *Let K be a number field and p be a prime. Then there exists a positive integer n depending on K and p such that for any elliptic curve E over K with no complex multiplication, we have*

$$\rho_{E,p}(\mathrm{G}_K) \supseteq 1 + p^n\mathrm{M}_2(\mathbb{Z}_p).$$

We will deduce Theorem 1.2 from the following more precise result of this paper in Proposition 2.10. We use the following conventions:

$$\begin{aligned} 1 + p^0\mathbb{Z}_p &:= \mathbb{Z}_p^\times, \\ 1 + p^0\mathrm{M}_2(\mathbb{Z}_p) &:= \mathrm{GL}_2(\mathbb{Z}_p), \\ 1 + p^0\mathrm{M}_2(\mathbb{Z}/p\mathbb{Z}) &:= \mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z}). \end{aligned}$$

Theorem 1.3. *For a prime p, there exists an integer $n \geq 0$ satisfying the following condition $(\boldsymbol{C})_p$.*
$(\boldsymbol{C})_p$*: Let K be a number field. Then there exists a finite subset $\Sigma \subseteq K$ depending on p such that for an elliptic curve E over K, the condition $j(E) \notin \Sigma$ implies*

$$\rho_{E,p}(\mathrm{G}_K) \supseteq (1 + p^n \mathrm{M}_2(\mathbb{Z}_p))^{\det=1}.$$

Let $n(p) \geq 0$ be the minimum integer n satisfying $(\boldsymbol{C})_p$. Then we have

$$n(p) \leq \begin{cases} 0 & \text{if } p \geq 23, \\ 1 & \text{if } p = 19,\ 17,\ 13,\ 11, \\ 2 & \text{if } p = 7, \\ 3 & \text{if } p = 5, \\ 5 & \text{if } p = 3, \\ 11 & \text{if } p = 2. \end{cases}$$

The estimate in Theorem 1.3 is best possible for $p \geq 7$.

Corollary 1.4. *We have*

$$n(p) = \begin{cases} 0 & \text{if } p \geq 23, \\ 1 & \text{if } p = 19,\ 17,\ 13,\ 11, \\ 2 & \text{if } p = 7. \end{cases}$$

Proof. Since $g(X_0(19)) = 1$, the integer $n(19)$ cannot be 0. We also have $g(X_0(17)) = 1$, $g(X_0(13)) = 0$, $g(X_0(11)) = 1$ and $g(X_0(7^2)) = 1$. Thus we get the result in a similar way.

□

Remark 1.5. *If $p \geq 23$ and*

$$\begin{cases} K \not\supseteq \text{(the quadratic subfield of } \mathbb{Q}(\zeta_p)) & \text{when } p \not\equiv \pm 3 \mod 8, \\ \text{there exists an inclusion } K \hookrightarrow \mathbb{Q}_p & \text{when } p \equiv \pm 3 \mod 8, \end{cases}$$

then the result of Theorem 1.3 follows from [3, Theorem 7; 21, Proposition 1.40, 1.43; 9, p.116-118].

If $p = 17$, 19 and $K = \mathbb{Q}$, the result of Theorem 1.3 follows from [3, Theorem 7; 8, Theorem (4.1); 9, p.116-118].

If $p = 2$ and $K = \mathbb{Q}$, the Galois image is studied in [14], assuming that an elliptic curve has no complex multiplication, and all its 2-torsion points are rational.

Remark 1.6. *There are many other studies of the Galois images associated to elliptic curves over number fields or of rational points on modular curves in [2,4,6,7,10-13,15,16,18,20]. Several questions related to the subject of this paper are raised in [19, p.187].*

The contents of this paper are as follows:

In Section 2, we deduce Theorem 1.2 from Theorem 1.3 by studying the determinants.

In Section 3, we regard elliptic curves as rational points on modular curves, and reduce Theorem 1.3 to a genus estimate. Replacing K by its finite extension, we may assume that K contains a primitive $p^{n(p)+1}$-st root $\zeta_{p^{n(p)+1}}$ of unity. Suppose an elliptic curve E/K does not satisfy $\rho_{E,p}(\mathrm{G}_K) \supseteq (1 + p^{n(p)} \mathrm{M}_2(\mathbb{Z}_p))^{\det=1}$. Then we have $\rho_{E,p}(\mathrm{G}_K) \mod p^{n(p)+1} \subseteq H$ for some subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{n(p)+1}\mathbb{Z})$ satisfying $H \not\supseteq (1+p^{n(p)}\mathrm{M}_2(\mathbb{Z}/p^{n(p)+1}\mathbb{Z}))^{\det=1}$. Thus E/K determines a rational point on the modular curve X_H corresponding to H. If the genus $g(X_H)$ of X_H is greater than or equal to 2, we conclude that X_H has only finitely many rational points by Mordell's conjecture ([3, Theorem 7]). Since there are only finitely many subgroups H as above, the number of the j-invariants of E/K not satisfying $\rho_{E,p}(\mathrm{G}_K) \supseteq (1+p^{n(p)}\mathrm{M}_2(\mathbb{Z}_p))^{\det=1}$ is finite. Thus Theorem 1.3 will follow.

In Section 4 - 7, we prove $g(X_H) \geq 2$. In section 4, we prepare for estimating $g(X_H)$. Put $G = \mathrm{SL}_2(\mathbb{Z}/p^{n(p)+1}\mathbb{Z})$, $\sigma = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, $\tau = \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$, $u = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. For $\alpha \in G$, let $\mathrm{Conj}(\alpha)$ be the conjugacy class containing α. If $H \ni -1$, we have

$$g(X_H) = 1 + \frac{1}{12}[G:H]\left(1 - 3\frac{\sharp H \cap \mathrm{Conj}(\sigma)}{\sharp \mathrm{Conj}(\sigma)} - 4\frac{\sharp H \cap \mathrm{Conj}(\tau)}{\sharp \mathrm{Conj}(\tau)} - 6\frac{\sharp \langle u \rangle \backslash G/H}{[G:H]}\right).$$

We calculate the number of elements conjugate to σ, τ, u contained in maximal subgroups of $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$.

In Section 5, we calculate the number of elements conjugate to σ, τ, u contained in $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$, and study the fiber of the mod p^m map $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \cap \mathrm{Conj}(\alpha) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z}) \cap \mathrm{Conj}(\alpha)$, where $1 \leq m \leq n$ and $\alpha = \sigma, \tau, u$. For integers $1 \leq m \leq n$, let $f_{n,m} : \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z})$ be the mod p^m map. If $m < n \leq 2m$ and $\alpha = \sigma, \tau, u$, we see that $\alpha^{-1}(f_{n,m}^{-1}(\alpha) \cap \mathrm{Conj}(\alpha))$ is a subgroup of $(1+p^m\mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z}))^{\det=1} \cong \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})^{\mathrm{Tr}=0}$ and isomorphic to $(\mathbb{Z}/p^{n-m}\mathbb{Z})^2$.

In Section 6, we control the number of elements conjugate to σ, τ, u contained in H, by combining the result of Section 5 with the property $H \not\supseteq (1+p^{n(p)}\mathrm{M}_2(\mathbb{Z}/p^{n(p)+1}\mathbb{Z}))^{\det=1}$.

In Section 7, we prove $g(X_H) \geq 2$ by using the results of Section 4, 5, 6.

I would like to thank my supervisor Professor Takeshi Saito for helpful advice and warm encouragement. This work was partly supported by 21st Century COE Program in The University of Tokyo, A Base for New Developments of Mathematics into Science and Technology.

2. Deduction of Theorem 1.2

In order to deduce Theorem 1.2 from Theorem 1.3, we need some facts in group theory.

Lemma 2.1. *Let p be a prime and H be a closed subgroup of $\mathrm{GL}_2(\mathbb{Z}_p)$. Then H contains $\mathrm{SL}_2(\mathbb{Z}_p)$ if and only if $H \bmod p^2$ contains $\mathrm{SL}_2(\mathbb{Z}/p^2\mathbb{Z})$. In particular, if a subgroup $H' \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ for $n \geq 3$ maps surjectively $\bmod p^2$ onto $\mathrm{SL}_2(\mathbb{Z}/p^2\mathbb{Z})$, then $H' = \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$.*

Assume $p \geq 5$. Then H contains $\mathrm{SL}_2(\mathbb{Z}_p)$ if and only if $H \bmod p$ contains $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$. In particular, if a subgroup $H' \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ for $n \geq 2$ maps surjectively $\bmod p$ onto $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$, then $H' = \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$.

Proof. See [17, IV-23].

□

Lemma 2.2. *Let $n \geq 1$ be an integer and let p be a prime. If $p = 2$, assume $n \geq 2$. Let H be a closed subgroup of $\mathrm{GL}_2(\mathbb{Z}_p)$. Then H contains $1+p^n\mathrm{M}_2(\mathbb{Z}_p)$ (resp. $(1+p^n\mathrm{M}_2(\mathbb{Z}_p))^{\det=1}$) if and only if $H \bmod p^{n+1}$ contains $1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$ (resp. $(1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))^{\det=1}$).*

Proof. The "only if" part is trivial.

Assume $H \bmod p^{n+1}$ contains $1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$. We will show that H contains $1+p^n\mathrm{M}_2(\mathbb{Z}_p)$. For each $s \in 1+p^n\mathrm{M}_2(\mathbb{Z}_p)$ and for any $a \geq 1$, we will find an element $x_a \in H$ with $x_a \equiv s \bmod p^{n+a}$. Then $s = (x_a \bmod p^{n+a})_{a\geq 1} \in H$, as desired. We show the existence of x_a's by induction on a. For $a = 1$, it follows from the assumption. Assume it is true for $a\,(\geq 1)$ i.e. $H \bmod p^{n+a} \supseteq 1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+a}\mathbb{Z})$. We first show that for $s \in 1+p^n\mathrm{M}_2(\mathbb{Z}_p)$ with $s \equiv 1 \bmod p^{n+a}$, there exists $x \in H$ with $x \equiv s \bmod p^{n+a+1}$. We write $s = 1+p^{n+a}u$ with some $u \in \mathrm{M}_2(\mathbb{Z}_p)$. By the hypothesis of induction, we find $y := 1+p^{n+a-1}u+p^{n+a}v \in H$ for some $v \in \mathrm{M}_2(\mathbb{Z}_p)$. Put $x := y^p$. Then $x \in H$ and $x = 1+p^{n+a}u+p^{n+a+1}\cdots \equiv s \bmod p^{n+a+1}$. For a general $s \in 1+p^n\mathrm{M}_2(\mathbb{Z}_p)$, the induction hypothesis shows that $(s \bmod p^{n+a})^{-1} \in H \bmod p^{n+a}$. Hence we have $st \equiv 1 \bmod p^{n+a}$ for some $t \in H$. By the above argument, we find $x \in H$ with $x \equiv st \bmod p^{n+a+1}$. Therefore $s \equiv t^{-1}x \bmod p^{n+a+1}$ and $t^{-1}x \in H$.

For the "det $= 1$" version, it follows from the same argument as in the previous lemma ([17, IV-23]).

□

Lemma 2.3. *Let $n \geq 1$ be an integer. Let H be a subgroup of $\mathrm{GL}_2(\mathbb{Z}_p)$ containing $1+p^n\mathrm{M}_2(\mathbb{Z}_p)$, and H' be a closed subgroup of $\mathrm{GL}_2(\mathbb{Z}_p)$ which is a subgroup of H of index 2. If $p \geq 3$, then $H' \supseteq 1+p^n\mathrm{M}_2(\mathbb{Z}_p)$; if $p = 2$ and $n \geq 2$, $H' \supseteq 1+p^{n+1}\mathrm{M}_2(\mathbb{Z}_p)$.*

Proof. For any $X \in \mathrm{M}_2(\mathbb{Z}_p)$, we have $H \ni 1+p^nX$ and $H' \ni (1+p^nX)^2 = 1+2p^nX+p^{2n}X^2$. This implies $H' \bmod p^{n+1} \supseteq 1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$ if $p \geq 3$ (*resp.* $H' \bmod p^{n+2} \supseteq 1+p^{n+1}\mathrm{M}_2(\mathbb{Z}/p^{n+2}\mathbb{Z})$ if $p = 2$ and $n \geq 2$). Since H' is closed, applying the previous lemma, we get the result.

□

Corollary 2.4. *Let $H \subseteq \mathrm{GL}_2(\mathbb{Z}_p)$ be a closed subgroup. If $\langle H, -1\rangle \supseteq 1+p^n\mathrm{M}_2(\mathbb{Z}_p)$ for some integer $n \geq 1$, then*

$$H \supseteq \begin{cases} 1+p^n\mathrm{M}_2(\mathbb{Z}_p) & \text{if } p \geq 3, \\ 1+p^{n+1}\mathrm{M}_2(\mathbb{Z}_p) & \text{if } p = 2 \text{ and } n \geq 2. \end{cases}$$

Lemma 2.5. *Let $n \geq 1$ be an integer and assume $p \geq 3$. Let H be a subgroup of $\mathrm{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$. If $\det(H) = (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times$, then*

$$\det|_{H\cap(1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))} : H \cap (1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})) \longrightarrow 1+p^n\mathbb{Z}/p^{n+1}\mathbb{Z}$$

is surjective.

Proof. We first show the surjectivity when $n = 1$. Let $x \in H$ be an element such that $\det x$ is a generator of $(\mathbb{Z}/p^2\mathbb{Z})^\times$. Put $\overline{x} := x \bmod p$ and $a := \mathrm{ord}\,\overline{x}$. We first show $(p,a) = 1$. Since $\det\overline{x}$ generates $\mathbb{F}_p^\times$, $\overline{x}$ is not a scalar. We consider the commutative subalgebra $\mathbb{F}_p[\overline{x}]$ of $\mathrm{M}_2(\mathbb{F}_p)$. If we put $t := \mathrm{Tr}\,\overline{x}$ and $d := \det\overline{x}$, then $\overline{x}^2 - t\overline{x} + dE = 0$ and $\mathbb{F}_p[\overline{x}] \cong \mathbb{F}_p[X]/(X^2-tX+d)$. X^2-tX+d does not have multiple roots because d generates $\mathbb{F}_p^\times$. Hence we have $\mathbb{F}_p[\overline{x}] \cong \mathbb{F}_p \times \mathbb{F}_p$ or $\mathbb{F}_{p^2}$. Therefore, $\overline{x} \in \mathbb{F}_p[\overline{x}]^\times \cong \mathbb{F}_p^\times \times \mathbb{F}_p^\times$ or $\mathbb{F}_{p^2}^\times$ and $(p,a) = 1$.

Since $\overline{x}^a = 1$, $x^a \in 1+p\mathrm{M}_2(\mathbb{Z}/p^2\mathbb{Z})$. Therefore, $\mathrm{ord}\,x^a = 1$ or p, and $\mathrm{ord}\,(\det x)^a = \mathrm{ord}\,(\det x^a) = 1$ or p. Thus, we have $\mathrm{ord}\,(\det x)^a = \mathrm{ord}\,(\det x^a) = p$ because $p | \mathrm{ord}\,(\det x)$ and $(p,a) = 1$. Hence, $\det x^a$ is a generator of $1+p\mathbb{Z}/p^2\mathbb{Z}$.

Next we show the surjectivity for general $n \geq 1$. Since $\det|_{H \bmod p^2} = (\det|_H) \bmod p^2$ is surjective, there exists an element $y \in (H \bmod p^2) \cap (1 + p\mathrm{M}_2(\mathbb{Z}/p^2\mathbb{Z}))$ such that $\det y$ is a generator of $1 + p\mathbb{Z}/p^2\mathbb{Z}$. We choose a lift $\tilde{y} \in H$ of y. Then $\tilde{y} \in 1 + p\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$. Taking the p^{n-1}-st power, we have $\tilde{y}^{p^{n-1}} \in H \cap (1 + p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))$. Since $\det y = \det \tilde{y} \bmod p^2$ generates $1 + p\mathbb{Z}/p^2\mathbb{Z}$, we can write $\det \tilde{y} = 1 + pz$ for some $z \in (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times$. Therefore $\det \tilde{y}^{p^{n-1}} = 1 + p^n z$ generates $1 + p^n\mathbb{Z}/p^{n+1}\mathbb{Z}$.

□

Lemma 2.6. *Take two integers $n > r \geq 1$. If $p = 2$, assume $r \geq 2$. Let H be a subgroup of $\mathrm{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$. If $\det(H)$ contains $1 + p^r\mathbb{Z}/p^{n+1}\mathbb{Z}$ and if H contains $(1 + p^{n-r}\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))^{\det=1}$, then*

$$\det|_{H\cap(1+p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))} : H \cap (1 + p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})) \longrightarrow 1 + p^n\mathbb{Z}/p^{n+1}\mathbb{Z}$$

is surjective. In particular, H contains $1 + p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$.

Proof. Let $x \in H$ be an element such that $\det x$ is a generator of $1 + p^r\mathbb{Z}/p^{n+1}\mathbb{Z}$. Put $y := x^{(p-1)^2(p+1)}$. We see that y is also a generator of $1 + p^r\mathbb{Z}/p^{n+1}\mathbb{Z}$ and y belongs to a p-Sylow subgroup of $\mathrm{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$, which is conjugate to $\left\{\begin{pmatrix} 1+p* & * \\ p* & 1+p* \end{pmatrix}\right\}$. We can write $y = 1+pz+\epsilon$, where ϵ is conjugate to $\begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$. Put $w := y^{p^{n-r}}$. We see that w belongs to $1+p^{n-r}\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$ and $\det w$ is a generator of $1+p^n\mathbb{Z}/p^{n+1}\mathbb{Z}$. Multiplying w by elements of $(1+p^{n-r}\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))^{\det=1}$, we get an element $h \in H \cap (1 + p^n\mathrm{M}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}))$ such that $\det h$ generates $1 + p^n\mathbb{Z}/p^{n+1}\mathbb{Z}$.

□

Corollary 2.7. *Let $H \subseteq \mathrm{GL}_2(\mathbb{Z}_p)$ be a closed subgroup and n, $r \geq 0$ be integers. Assume $r \geq 2$ if $p = 2$. If $H \supseteq (1 + p^n\mathrm{M}_2(\mathbb{Z}_p))^{\det=1}$ and if $\det(H) \supseteq 1 + p^r\mathbb{Z}_p$, then $H \supseteq 1 + p^{n+r}\mathrm{M}_2(\mathbb{Z}_p)$.*

Proof. By Lemma 2.2, it suffices to show the result mod p^{n+r+1}. If $n = 0$, it is trivial. Assume $n \geq 1$. If $r = 0$, it follows from Lemma 2.5. If $r \geq 1$, it follows from Lemma 2.6.

□

As a consequence of Theorem 1.3, we get the following.

Theorem 2.8. *Let K be a number field, p be a prime. Let $n(p)$, Σ be as in Theorem 1.3. Suppose that the image of the p-adic cyclotomic character $\chi_p : \mathrm{G}_K \longrightarrow \mathbb{Z}_p^\times$ contains $1 + p^r\mathbb{Z}_p$ with $r \geq 0$ an integer. Assume $r \geq 2$ if $p = 2$. Put*

$$n = r + n(p).$$

Then for an elliptic curve E over K, the condition $j(E) \notin \Sigma$ implies $\rho_{E,p}(\mathrm{G}_K) \supseteq 1 + p^n\mathrm{M}_2(\mathbb{Z}_p)$.

Proof. It follows from Theorem 1.3 and Corollary 2.7.

□

We show that there exists a lower bound of the images of $\rho_{E,p}$ if we take E's having only finitely many j-invariants.

Lemma 2.9. *Let K be a number field. Fix an element $j \in K$. Assume that an elliptic curve E over K with j-invariant j has no complex multiplication. Take a prime p. Then there exists a positive integer n depending on p and j such that for any elliptic curve E over K with j-invariant j, we have $\rho_{E,p}(\mathrm{G}_K) \supseteq 1+p^n\mathrm{M}_2(\mathbb{Z}_p)$.*

Proof. Take an E/K with j-invariant j. By Theorem 1.1, we have $\rho_{E,p}(\mathrm{G}_K) \supseteq 1+p^{n_0}\mathrm{M}_2(\mathbb{Z}_p)$ for some n_0. Let E'/K have j-invariant j. Since E has no complex multiplication, we have $j \neq 0, 1728$. Hence there exists a quadratic extension L of K satisfying $E \otimes_K L \cong E' \otimes_K L$ (see [22, p.308]). Therefore $\rho_{E,p}(\mathrm{G}_L)$ is conjugate to $\rho_{E',p}(\mathrm{G}_L)$. Since $[\rho_{E,p}(\mathrm{G}_K) : \rho_{E,p}(\mathrm{G}_L)]$ is 1 or 2, applying Lemma 2.3, we get the result.

□

Now we can deduce Theorem 1.2 from Theorem 1.3 and some facts in group theory.

Proposition 2.10. *Theorem 1.3 and Lemma 2.9 imply Theorem 1.2.*

Proof. Theorem 1.3 implies Theorem 2.8. Since Σ is a finite set, elliptic curves E over K with $j(E) \in \Sigma$ have bounded Galois images by Lemma 2.9.

□

3. Modular curves

We regard an elliptic curve with a specific Galois image as a rational point on a certain modular curve, and reduce Theorem 1.3 to a genus estimate.

Now we give a brief review of modular curves. For more details, see [1,5]. Let N be a positive integer and k be a field of characteristic 0. For an elliptic curve E over k and an integer $N \geq 1$, let $E[N] = \mathrm{Ker}\,([N] : E \longrightarrow E)$ be the kernel of multiplication by N on E, and let $\overline{\rho}_{E,N} : \mathrm{G}_k \longrightarrow \mathrm{Aut}(E[N](\overline{k})) \cong \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ be the mod N representation determined by the action of G_k on $E[N](\overline{k})$. A level N-structure on E is an isomorphism

$$\gamma : (\mathbb{Z}/N\mathbb{Z})^2 \longrightarrow E[N].$$

Let $Y(N) \longrightarrow \mathrm{Spec}(\mathbb{Q}(\zeta_N))$ be the moduli of elliptic curves with level N-structure. We have a right action of $G = \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ on $Y(N)$ over $\mathbb{Q}(\zeta_N)$:

$$[E, \gamma] \mapsto [E, \gamma \circ h]$$

where E is an elliptic curve over k, γ a level N-structure on E and $h \in G$.

For a subgroup $H \subseteq G$, put Y_H to be the quotient $Y(N)/H$. The quotient $Y_H \longrightarrow \mathrm{Spec}(\mathbb{Q}(\zeta_N))$ is an affine smooth curve. Let E be an elliptic curve over k. Choose a basis $\langle \epsilon_1, \epsilon_2 \rangle$ of $E[N](\overline{k})$. Then the pair $(E, \langle \epsilon_1, \epsilon_2 \rangle)$ defines an element P of $Y(N)(\overline{k})$. Let $Q \in Y_H(\overline{k})$ be the image of P via the map $Y(N)(\overline{k}) \longrightarrow Y_H(\overline{k})$ induced by the natural map $Y(N) \longrightarrow Y_H$. If $\overline{\rho}_{E,N}(\mathrm{G}_k) \subseteq H$ with respect to $\langle \epsilon_1, \epsilon_2 \rangle$, then Q lies in $Y_H(k)$. Note that if two subgroups $H, H' \subseteq G$ are $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$-conjugate, then Y_H and $Y_{H'}$ are conjugate.

Lemma 3.1. *Let k be a field of characteristic 0. If $Y_H(k)$ is finite, then there exists a finite set $\Sigma \subseteq k$ satisfying the following condition:*

For an elliptic curve E over k, if a conjugate of $\overline{\rho}_{E,N}(\mathrm{G}_k)$ is contained in H, then $j(E) \in \Sigma$.

Let X_H be the smooth compactification of Y_H. The following is the famous theorem known as Mordell's conjecture proved by Faltings. It shows that a curve X over a number field has only finitely many rational points if its genus $g(X)$ is greater than or equal to 2.

Theorem 3.2. *([3, Theorem 7]) Let K be a number field and X/K be a proper smooth curve. If $g(X) \geq 2$, then $X(K)$ is finite.*

Now we compute the genus of X_H explicitly. As in Section 1, put

$$\sigma := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},\ \tau := \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix},\ u := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}).$$

We have $\operatorname{ord}\sigma = 4$ and $\operatorname{ord}\tau = 6$. Note that $\sigma^{-1} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\tau^{-1} = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$. For $\alpha \in \mathrm{SL}_2(\mathbb{Z})$, we also use the same letter to denote the reduction of α. Put

$$\mathrm{Fix}_\alpha := \{gH \in G/H | \alpha gH = gH\}$$

and

$$g_H := 1 + \frac{1}{12}\sharp G/H - \frac{1}{4}\sharp \mathrm{Fix}_\sigma - \frac{1}{3}\sharp \mathrm{Fix}_\tau - \frac{1}{2}\sharp I\backslash G/H.$$

Here I is the cyclic subgroup of G generated by $u = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Proposition 3.3. *([21, Proposition 1.40]) Let H be a subgroup of $G = \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$. Assume that H contains -1. Then the genus $g(X_H)$ of the modular curve X_H is given by*

$$g(X_H) = g_H.$$

Let p be a prime and consider subgroups of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$. A Borel subgroup is a subgroup which is conjugate to $\left\{\begin{pmatrix} * & * \\ 0 & * \end{pmatrix}\right\}$; the normalizer of a split Cartan subgroup is conjugate to $\left\{\begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}, \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix}\right\}$. When $p \geq 3$, the normalizer of a non-split Cartan subgroup is conjugate to $\left\{\begin{pmatrix} x & y \\ \lambda y & x \end{pmatrix}, \begin{pmatrix} x & y \\ -\lambda y & -x \end{pmatrix} |(x,y) \in \mathbb{F}_p \times \mathbb{F}_p \setminus \{(0,0)\}\right\}$, where $\lambda \in \mathbb{F}_p^\times \setminus (\mathbb{F}_p^\times)^2$ is a fixed element. Assume $p \geq 5$. The quotient group $\mathrm{PGL}_2(\mathbb{Z}/p\mathbb{Z})$ of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$ has a subgroup which is isomorphic to S_4; it has a subgroup which is isomorphic to A_5 if and only if $p \equiv 0, \pm 1 \mod 5$ ([18, p.281]). Take a subgroup H (of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$) whose order is prime to p. We call H an exceptional subgroup if it is the inverse image of a subgroup which is isomorphic to A_4, S_4 or A_5 by the natural surjection $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z}) \longrightarrow \mathrm{PGL}_2(\mathbb{Z}/p\mathbb{Z})$.

Proposition 3.4. *([18, p.284]) Let $p \geq 3$ be a prime and H be a subgroup of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$. If p divides the order of H, then H contains $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ or H is contained in a Borel subgroup of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$. If p does not divide the order of H, then H is contained in the normalizer of a (split or non-split) Cartan subgroup of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$ or an exceptional subgroup of $\mathrm{GL}_2(\mathbb{Z}/p\mathbb{Z})$.*

Put

$$\begin{aligned}
B &:= \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \right\} \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}), \\
C &:= \left\{ \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}, \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix} \right\} \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}), \\
D &:= \left\{ \begin{pmatrix} x & y \\ \lambda y & x \end{pmatrix}, \begin{pmatrix} x & y \\ -\lambda y & -x \end{pmatrix} \right\} \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}), \\
E &:= (\text{an exceptional subgroup}) \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}).
\end{aligned}$$

From now on, we use the letter E to denote this subgroup, not meaning an elliptic curve. The genera of the modular curves corresponding to B, C, D are known as follows.

Proposition 3.5. *([21, Proposition 1.40, 1.43; 9, p.117]) Let $N = p \geq 5$ be a prime. We have*

$$\begin{aligned}
g_B &= \frac{1}{12}(p - 6 - 3\left(\frac{-1}{p}\right) - 4\left(\frac{-3}{p}\right)); \\
g_C &= \frac{1}{24}(p^2 - 8p + 11 - 4\left(\frac{-3}{p}\right)); \\
g_D &= \frac{1}{24}(p^2 - 10p + 23 + 6\left(\frac{-1}{p}\right) + 4\left(\frac{-3}{p}\right)).
\end{aligned}$$

We have $g_B \geq 2$ if and only if $p \geq 23$; we have $g_C \geq 2$ if and only if $p \geq 11$; we have $g_D \geq 2$ if and only if $p \geq 13$.

Remark 3.6. *We can also calculate these genera by using Lemma 4.9.*

Put

$$\delta_H := [G : H]^{-1}(\sharp G/H - 3\sharp \mathrm{Fix}_\sigma - 4\sharp \mathrm{Fix}_\tau - 6\sharp I\backslash G/H)$$

so that

$$g_H = 1 + \frac{1}{12}[G : H]\delta_H.$$

As g_H is an integer, we have $g_H \geq 2$ if and only if $\delta_H > 0$.

Definition 3.7. *Let $n \geq 1$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ be a subgroup. We call H a slim subgroup if*

$$H \not\supseteq (1 + p^{n-1}\mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z}))^{\det=1}.$$

In this definition, notice that if $n = 1$, then a slim subgroup is just a proper subgroup.

In order to prove Theorem 1.3, it suffices to estimate δ_H for any slim subgroup H.

Proposition 3.8. *If $\delta_H > 0$ for any slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{n'(p)+1}\mathbb{Z})$, then Theorem 1.3 holds. Here we put*

$$n'(p) := \begin{cases} 0 & \textit{if } p \geq 23, \\ 1 & \textit{if } p = 19, 17, 13, 11, \\ 2 & \textit{if } p = 7, \\ 3 & \textit{if } p = 5, \\ 5 & \textit{if } p = 3, \\ 10 & \textit{if } p = 2. \end{cases}$$

Proof. Put

$$\xi := \begin{cases} 0 & \text{if } p \geq 3, \\ 1 & \text{if } p = 2. \end{cases}$$

Let E be an elliptic curve over K satisfying $\rho_{E,p}(\mathrm{G}_K) \not\supseteq (1 + p^{n'(p)+\xi}\mathrm{M}_2(\mathbb{Z}_p))^{\det=1}$. We show that $j(E) \in K$ takes only finitely many values. Replacing K by $K(\zeta_{p^{n'(p)+1}})$, we may assume $\overline{\rho}_{E,p^{n'(p)+1}}(\mathrm{G}_K) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{n'(p)+1}\mathbb{Z})$. We may also assume that $\overline{\rho}_{E,p^{n'(p)+1}}(\mathrm{G}_K)$ is contained in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{n'(p)+1}\mathbb{Z})$ satisfying $H \ni -1$. To see this, we consider two cases ($n'(p) = 0$ or $n'(p) \geq 1$). When $n'(p) = 0$ (equivalently $p \geq 23$), we have $\overline{\rho}_{E,p}(\mathrm{G}_K) \subsetneq \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ by Lemma 2.1, thus $\overline{\rho}_{E,p}(\mathrm{G}_K) \subseteq B, C, D, E$ by Proposition 3.4. But B, C, D, E contains -1. When $n'(p) \geq 1$, Corollary 2.4 shows $\langle \rho_{E,p}(\mathrm{G}_K), -1 \rangle \not\supseteq (1 + p^{n'(p)}\mathrm{M}_2(\mathbb{Z}_p))^{\det=1}$. Thus $H = \langle \rho_{E,p}(\mathrm{G}_K), -1 \rangle \bmod p^{n'(p)+1}$ is a slim subgroup by Lemma 2.2.

By the hypothesis and Proposition 3.3, we have $g(X_H) = g_H \geq 2$. By Theorem 3.2, we see that $X_H(K)$ is finite, hence $Y_H(K)$ is also finite. Since there are only finitely many subgroups H as above, Theorem 1.3 follows from Lemma 3.1.

□

We prove $\delta_H > 0$ for any subgroup H as in Proposition 3.8. More explicitly, we prove the following theorem in Section 7.

Theorem 3.9. *1. For a subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$, we have $\delta_H > 0$ if one of the following conditions is satisfied.*

- $H \subseteq B$ *and* $p \geq 23$
- $H \subseteq C$ *and* $p \geq 11$
- $H \subseteq D$ *and* $p \geq 13$
- $H \subseteq E$ *and* $p \geq 17$

2. For a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^2\mathbb{Z})$, we have $\delta_H > 0$ if one of the following conditions is satisfied.

- $H \bmod p \subseteq B$ *and* $p \geq 11$
- $H \bmod p \subseteq D$ *and* $p \geq 11$
- $H \bmod p \subseteq E$ *and* $p \geq 11$

3. For a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^3\mathbb{Z})$, we have $\delta_H > 0$ if one of the following conditions is satisfied.

- $H \bmod p \subseteq B$ *and* $p \geq 7$
- $H \bmod p \subseteq C$ *and* $p \geq 5$
- $H \bmod p \subseteq D$ *and* $p \geq 7$
- $H \bmod p \subseteq E$ *and* $p \geq 7$

4. For a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/5^4\mathbb{Z})$*, we have* $\delta_H > 0$ *if one of the following conditions is satisfied.*

- $H \bmod 5 \subseteq B$
- $H \bmod 5 \subseteq D$
- $H \bmod 5 \subseteq E$

5. For a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/3^6\mathbb{Z})$*, we have* $\delta_H > 0$ *if one of the following conditions is satisfied.*

- $H \bmod 3 \subseteq B$
- $H \bmod 3 \subseteq D$
- $H \bmod 3 \subseteq E$
- $H \bmod 3 = \mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$

6. For a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{10}\mathbb{Z})$*, we have* $\delta_H > 0$ *if one of the following conditions is satisfied.*

- $H \bmod 2 \subseteq$ (*subgroup of order* 3)
- $H \bmod 2 = \mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$

7. For a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{11}\mathbb{Z})$*, we have* $\delta_H > 0$ *if the following condition is satisfied.*

- $H \bmod 2 \subseteq B$

4. Calculation of conjugate elements in $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$

We rewrite the value δ_H in terms of the number of conjugate elements.

Lemma 4.1. *Let* G *be a finite group and* H *be a subgroup of* G*. Take an element* a *of* G*. Then*

$$\sharp\mathrm{Fix}_a/[G:H] = \sharp(H \cap \mathrm{Conj}(a))/\sharp\mathrm{Conj}(a).$$

Here we put $\mathrm{Conj}(a) := \{g^{-1}ag \in G | g \in G\}$*.*

Proof. Put $\widetilde{\mathrm{Fix}_a} := \{g \in G | gH = agH\}$. Then H acts on $\widetilde{\mathrm{Fix}_a}$ by $g \mapsto gh$ ($g \in \widetilde{\mathrm{Fix}_a},\ h \in H$) and we have $\widetilde{\mathrm{Fix}_a}/H = \mathrm{Fix}_a$. Hence, we get $\sharp\widetilde{\mathrm{Fix}_a}/\sharp H = \sharp\mathrm{Fix}_a$. The natural surjection $\widetilde{\mathrm{Fix}_a} \longrightarrow H \cap \mathrm{Conj}(a) : g \longmapsto g^{-1}ag$ induces a bijection $\mathrm{Z}(a)\backslash\widetilde{\mathrm{Fix}_a} \cong H \cap \mathrm{Conj}(a)$, where $\mathrm{Z}(a)$ is the centralizer of a in G, acting on $\widetilde{\mathrm{Fix}_a}$ by $g \mapsto zg$ ($g \in \widetilde{\mathrm{Fix}_a},\ z \in \mathrm{Z}(a)$). Hence $\sharp\widetilde{\mathrm{Fix}_a}/\sharp\mathrm{Z}(a) = \sharp(H \cap \mathrm{Conj}(a))$. If we take H to be G, we have $\sharp G/\sharp\mathrm{Z}(a) = \sharp\mathrm{Conj}(a)$. Combining these three equalities, we get the desired formula. □

For an integer $N \geq 1$, take a subgroup $H \subseteq G = \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$. By Lemma 4.1, we have

$$\delta_H = 1 - 3\frac{\sharp H \cap \mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)} - 4\frac{\sharp H \cap \mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} - 6\frac{\sharp I\backslash G/H}{[G:H]}. \tag{4.1}$$

Recall $\mathrm{Conj}(\alpha) = \{g^{-1}\alpha g \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z}) | g \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})\}$ for $\alpha \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$.

Assume $n \geq 1$ and $N = p^n$. For $0 \leq s \leq n$, we have

$$\mathrm{Fix}_{u^{p^s}} = \{gH \in G/H | \sharp I(gH) \leq p^s\},$$

where $I(gH)$ means the I-orbit of gH in G/H. We have

$$G/H = \mathrm{Fix}_{u^{p^n}} = \mathrm{Fix}_u \amalg \coprod_{s=0}^{n-1} (\mathrm{Fix}_{u^{p^{s+1}}} \setminus \mathrm{Fix}_{u^{p^s}})$$

and

$$\sharp I\backslash G/H = \sharp\mathrm{Fix}_u + \sum_{s=0}^{n-1}\frac{1}{p^{s+1}}(\sharp\mathrm{Fix}_{u^{p^{s+1}}} - \sharp\mathrm{Fix}_{u^{p^s}}) = \sum_{s=0}^{n-1}\frac{p-1}{p^{s+1}}\sharp\mathrm{Fix}_{u^{p^s}} + \frac{1}{p^n}[G:H].$$

By Lemma 4.1, we get

$$\frac{\sharp I\backslash G/H}{[G:H]} = \sum_{s=0}^{n-1}\frac{p-1}{p^{s+1}}\frac{\sharp H\cap \mathrm{Conj}(u^{p^s})}{\sharp\mathrm{Conj}(u^{p^s})} + \frac{1}{p^n}. \tag{4.2}$$

In particular, we have

$$\frac{\sharp I\backslash G/H}{[G:H]} \leq \sum_{s=0}^{t-1}\frac{p-1}{p^{s+1}}\frac{\sharp H\cap \mathrm{Conj}(u^{p^s})}{\sharp\mathrm{Conj}(u^{p^s})} + \frac{1}{p^t} \tag{4.3}$$

for $1 \leq t \leq n$.

When $p = 2$ or 3, we study the conjugacy classes of the maximal subgroups of $\mathrm{SL}_2(\mathbb{Z}/p^2\mathbb{Z})$.

Lemma 4.2. *We have*

$$\mathrm{SL}_2(\mathbb{Z}) = \langle\sigma,\tau\rangle = \langle\sigma,u\rangle = \langle\tau,u\rangle = \langle u,\, {}^t u\rangle.$$

Proof. The equality $\mathrm{SL}_2(\mathbb{Z}) = \langle\sigma,u\rangle$ is well-known ([21, p.16]). As $\sigma = u\tau$ and $({}^t u)^{-1}u\cdot({}^t u)^{-1} = \sigma$, we have the other equalities.

□

Lemma 4.3. *([21, Lemma 1.38]) For any positive integer N, the map*

$$\bmod N : \mathrm{SL}_2(\mathbb{Z}) \longrightarrow SL_2(\mathbb{Z}/N\mathbb{Z})$$

is surjective.

Lemma 4.4. *Let $N \geq 2$ be an integer and $A \subsetneq \mathrm{SL}_2(\mathbb{Z}/N^2\mathbb{Z})$ be a proper subgroup. Assume A maps surjectively* $\bmod N$ *onto* $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$*. Then A has no element which is* $\mathrm{GL}_2(\mathbb{Z}/N^2\mathbb{Z})$*-conjugate to u.*

Proof. Suppose A has an element which is $\mathrm{GL}_2(\mathbb{Z}/N^2\mathbb{Z})$-conjugate to u. Replacing A by its $\mathrm{GL}_2(\mathbb{Z}/N^2\mathbb{Z})$-conjugate, we may assume $A \ni u = \begin{pmatrix}1 & 1\\ 0 & 1\end{pmatrix}$. By the hypothesis, we see that A contains an element X satisfying $X \mod N = \begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}$. Since X has determinant 1, we can write $X = \begin{pmatrix}Na & 1+Nb\\ -1+Nb & Nd\end{pmatrix}$. We have

$$A \ni \begin{pmatrix}1 & Na\\ 0 & 1\end{pmatrix}\begin{pmatrix}Na & 1+Nb\\ -1+Nb & Nd\end{pmatrix}\begin{pmatrix}1 & Nd\\ 0 & 1\end{pmatrix} = \begin{pmatrix}0 & 1+Nb\\ -1+Nb & 0\end{pmatrix}.$$

Hence

$$A \ni \begin{pmatrix}0 & 1+Nb\\ -1+Nb & 0\end{pmatrix}^{-1}\begin{pmatrix}1 & 1\\ 0 & 1\end{pmatrix}\begin{pmatrix}0 & 1+Nb\\ -1+Nb & 0\end{pmatrix} = \begin{pmatrix}1 & 0\\ -1+2Nb & 1\end{pmatrix}.$$

Take an integer $\widetilde{b}$ such that $\widetilde{b} \mod N^2 = b$, and we get $A \ni \begin{pmatrix}1 & 0\\ -1+2Nb & 1\end{pmatrix}^{-1-2N\widetilde{b}} = \begin{pmatrix}1 & 0\\ 1 & 1\end{pmatrix}$. Then $A \supseteq \langle u,\, {}^t u\rangle = \mathrm{SL}_2(\mathbb{Z}/N^2\mathbb{Z})$, a contradiction.

□

Remark 4.5. *If $N = p \geq 5$ is a prime, there is no subgroup A as in the previous lemma by Lemma 2.1.*

Lemma 4.6. *The conjugacy classes of* $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ *are the following:*

$$\begin{aligned}
\mathrm{Conj}(1) =& \{1\},\\
\mathrm{Conj}(-1) =& \{-1\},\\
\mathrm{Conj}(\sigma) =& \left\{\sigma, \begin{pmatrix}1 & 2\\ -1 & -1\end{pmatrix}, \begin{pmatrix}2 & 1\\ -1 & 2\end{pmatrix}, \begin{pmatrix}-1 & 2\\ -1 & 1\end{pmatrix}, \begin{pmatrix}1 & 1\\ 2 & -1\end{pmatrix}, \begin{pmatrix}-1 & 1\\ 2 & 1\end{pmatrix}\right\},\\
\mathrm{Conj}(-\sigma) =& -\mathrm{Conj}(\sigma),\\
\mathrm{Conj}(\tau) =& \left\{\tau, \begin{pmatrix}1 & -1\\ 1 & 0\end{pmatrix}, \begin{pmatrix}0 & 1\\ -1 & 1\end{pmatrix}, \begin{pmatrix}0 & -1\\ 1 & 1\end{pmatrix}, \begin{pmatrix}-1 & 1\\ 1 & 2\end{pmatrix}, \begin{pmatrix}-1 & -1\\ -1 & 2\end{pmatrix},\right.\\
& \left.\begin{pmatrix}2 & 1\\ 1 & -1\end{pmatrix}, \begin{pmatrix}2 & -1\\ -1 & -1\end{pmatrix}\right\},\\
\mathrm{Conj}(-\tau) =& -\mathrm{Conj}(\tau),\\
\mathrm{Conj}(u) =& \left\{u, \begin{pmatrix}1 & 0\\ -1 & 1\end{pmatrix}, \begin{pmatrix}2 & 1\\ -1 & 0\end{pmatrix}, \begin{pmatrix}-1 & 0\\ -1 & -1\end{pmatrix}, \begin{pmatrix}0 & 1\\ -1 & 2\end{pmatrix}, \begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\right\},\\
\mathrm{Conj}(-u) =& -\mathrm{Conj}(u),\\
\mathrm{Conj}(u^2) =& \left\{u^2, \begin{pmatrix}1 & 0\\ 2 & 1\end{pmatrix}, \begin{pmatrix}-1 & 2\\ 2 & -1\end{pmatrix}\right\},\\
\mathrm{Conj}(-u^2) =& -\mathrm{Conj}(u^2).
\end{aligned}$$

Now we determine maximal subgroups of $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ whose image mod 2 is $\mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$.

Lemma 4.7. *Let $A \subsetneq \mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ be a proper subgroup. Assume A maps surjectively* mod 2 *onto* $\mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$. *Then A is conjugate to*

$$A_1 := \langle \sigma, \begin{pmatrix}1 & 1\\ 2 & -1\end{pmatrix}\rangle,$$

which is a maximal subgroup, and is not a normal subgroup. We have $\langle A_1, u^2\rangle = \mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ and $A_1\backslash\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z}) = \{1, u, u^2, u^{-1}\}$.

Proof. By the hypothesis, we see that A contains a lift of $u \in \mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$. We have

$$\{X \in \mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z}) | X \bmod 2 = u\} = \{u, u^{-1}, -u, -u^{-1}, \alpha, \beta, \alpha^{-1}, \beta^{-1}\},$$

where $\alpha = \begin{pmatrix}1 & 1\\ 2 & -1\end{pmatrix}$ and $\beta = \begin{pmatrix}-1 & 1\\ 2 & 1\end{pmatrix}$. The first four elements are $\mathrm{GL}_2(\mathbb{Z}/4\mathbb{Z})$-conjugate to u. The next two elements are $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$-conjugate to σ and the last two to σ^{-1}. Since A has no element which is $\mathrm{GL}_2(\mathbb{Z}/4\mathbb{Z})$-conjugate to u, we have $A \cap \mathrm{Conj}(\sigma) \neq \emptyset$. Replacing A by its conjugate, we may assume $A \ni \sigma$. Repeating the same argument, we see that A contains at least one of $\alpha, \beta, \alpha^{-1}, \beta^{-1}$. We have $A_1 = \langle \sigma, \alpha\rangle = \langle \sigma, \alpha^{-1}\rangle$. Put $A_2 := \langle \sigma, \beta\rangle = \langle \sigma, \beta^{-1}\rangle$. By calculation, we have

$$\begin{aligned}
A_1 =& \left\{\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}, \begin{pmatrix}-1 & 0\\ 0 & -1\end{pmatrix}, \begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}, \begin{pmatrix}0 & -1\\ 1 & 0\end{pmatrix}, \begin{pmatrix}2 & 1\\ 1 & 1\end{pmatrix}, \begin{pmatrix}1 & -1\\ -1 & 2\end{pmatrix},\right.\\
& \left.\begin{pmatrix}2 & -1\\ -1 & -1\end{pmatrix}, \begin{pmatrix}-1 & 1\\ 1 & 2\end{pmatrix}, \begin{pmatrix}-1 & 2\\ -1 & 1\end{pmatrix}, \begin{pmatrix}1 & 2\\ 1 & -1\end{pmatrix}, \begin{pmatrix}1 & 1\\ 2 & -1\end{pmatrix}, \begin{pmatrix}-1 & -1\\ 2 & 1\end{pmatrix}\right\}
\end{aligned}$$

and

$$\begin{aligned}A_2 = \Big\{ &\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}, \begin{pmatrix}-1 & 0\\ 0 & -1\end{pmatrix}, \begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}, \begin{pmatrix}0 & -1\\ 1 & 0\end{pmatrix}, \begin{pmatrix}2 & -1\\ -1 & 1\end{pmatrix}, \begin{pmatrix}1 & 1\\ 1 & 2\end{pmatrix},\\ &\begin{pmatrix}2 & 1\\ 1 & -1\end{pmatrix}, \begin{pmatrix}-1 & -1\\ -1 & 2\end{pmatrix}, \begin{pmatrix}-1 & 2\\ 1 & 1\end{pmatrix}, \begin{pmatrix}1 & 2\\ -1 & -1\end{pmatrix}, \begin{pmatrix}1 & -1\\ 2 & -1\end{pmatrix}, \begin{pmatrix}-1 & 1\\ 2 & 1\end{pmatrix} \Big\}.\end{aligned}$$

The equalities $u^{-1}\sigma u = \begin{pmatrix}1 & 2\\ -1 & -1\end{pmatrix} \in A_2$ and $u^{-1}\alpha u = \begin{pmatrix}-1 & 1\\ 2 & 1\end{pmatrix} \in A_2$ show $A_2 = u^{-1}A_1u$. We also have $\langle A_1, u^2\rangle = \mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ by calculation.

□

Remark 4.8. *In the above lemma, all the subgroups of* $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ *conjugate to* A_1 *are* A_1, $u^{-1}A_1u$, $u^{-2}A_1u^2$ *and* uA_1u^{-1}.

Now we calculate the number of elements conjugate to σ, τ, u in the maximal subgroups B, C, D, E introduced in Section 3.

Lemma 4.9. *In* $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$, *the number of the elements conjugate to* σ, τ, u *in* B, C, D, E *are as follows.*

1.

$$B \cap \mathrm{Conj}(\sigma) = \begin{cases} \emptyset & \text{if } p \equiv -1 \mod 4,\\ \left\{ \begin{pmatrix}\sqrt{-1} & *\\ 0 & -\sqrt{-1}\end{pmatrix}, \begin{pmatrix}-\sqrt{-1} & *\\ 0 & \sqrt{-1}\end{pmatrix} \right\} & \text{if } p \equiv 1 \mod 4,\\ \left\{ \begin{pmatrix}1 & 1\\ 0 & 1\end{pmatrix} \right\} & \text{if } p = 2.\end{cases}$$

$$B \cap \mathrm{Conj}(\tau) = \begin{cases} \emptyset & \text{if } p \equiv -1 \mod 3,\\ \left\{ \begin{pmatrix}-\zeta_3 & *\\ 0 & -\zeta_3^{-1}\end{pmatrix}, \begin{pmatrix}-\zeta_3^{-1} & *\\ 0 & -\zeta_3\end{pmatrix} \right\} & \text{if } p \equiv 1 \mod 3,\\ \left\{ \begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix} \right\} & \text{if } p = 3.\end{cases}$$

$$B \cap \mathrm{Conj}(u) = \left\{ \begin{pmatrix}1 & s^2\\ 0 & 1\end{pmatrix} |(p, s) = 1 \right\}.$$

Hence we have

$$\sharp B \cap \mathrm{Conj}(\sigma) = \begin{cases} 0 & \text{if } p \equiv -1 \mod 4,\\ 2p & \text{if } p \equiv 1 \mod 4,\\ 1 & \text{if } p = 2.\end{cases}$$

$$\sharp B \cap \mathrm{Conj}(\tau) = \begin{cases} 0 & \text{if } p \equiv -1 \mod 3,\\ 2p & \text{if } p \equiv 1 \mod 3,\\ 1 & \text{if } p = 3.\end{cases}$$

$$\sharp B \cap \mathrm{Conj}(u) = \begin{cases} \frac{1}{2}(p-1) & \text{if } p \geq 3,\\ 1 & \text{if } p = 2.\end{cases}$$

2.

$$C \cap \mathrm{Conj}(\sigma) = \begin{cases} \left\{ \begin{pmatrix} 0 & x \\ -x^{-1} & 0 \end{pmatrix} \middle| (p,x)=1 \right\} \\ \text{if } p \equiv -1 \mod 4, \\ \left\{ \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix}, \begin{pmatrix} -\sqrt{-1} & 0 \\ 0 & \sqrt{-1} \end{pmatrix}, \begin{pmatrix} 0 & x \\ -x^{-1} & 0 \end{pmatrix} \middle| (p,x)=1 \right\} \\ \text{if } p \equiv 1 \mod 4, \\ \left\{ \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\} \\ \text{if } p = 2. \end{cases}$$

$$C \cap \mathrm{Conj}(\tau) = \begin{cases} \emptyset & \text{if } p \not\equiv 1 \mod 3, \\ \left\{ \begin{pmatrix} -\zeta_3 & 0 \\ 0 & -\zeta_3^{-1} \end{pmatrix}, \begin{pmatrix} -\zeta_3^{-1} & 0 \\ 0 & -\zeta_3 \end{pmatrix} \right\} & \text{if } p \equiv 1 \mod 3. \end{cases}$$

$$C \cap \mathrm{Conj}(u) = \begin{cases} \emptyset & \text{if } p \geq 3, \\ \left\{ \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\} & \text{if } p = 2. \end{cases}$$

Hence we have

$$\sharp C \cap \mathrm{Conj}(\sigma) = \begin{cases} p-1 & \text{if } p \equiv -1 \mod 4, \\ p+1 & \text{if } p \equiv 1 \mod 4, \\ 1 & \text{if } p = 2. \end{cases}$$

$$\sharp C \cap \mathrm{Conj}(\tau) = \begin{cases} 0 & \text{if } p \not\equiv 1 \mod 3, \\ 2 & \text{if } p \equiv 1 \mod 3. \end{cases}$$

$$\sharp C \cap \mathrm{Conj}(u) = \begin{cases} 0 & \text{if } p \geq 3, \\ 1 & \text{if } p = 2. \end{cases}$$

3.

$$D \cap \mathrm{Conj}(\tau) = \begin{cases} \left\{ \begin{pmatrix} 1/2 & 1/2 \cdot \sqrt{-3/\lambda} \\ 1/2 \cdot \sqrt{-3\lambda} & 1/2 \end{pmatrix}, \begin{pmatrix} 1/2 & -1/2 \cdot \sqrt{-3/\lambda} \\ -1/2 \cdot \sqrt{-3\lambda} & 1/2 \end{pmatrix} \right\} \\ \text{if } p \geq 5 \text{ and } p \equiv -1 \mod 3, \\ \emptyset \\ \text{if } p = 3 \text{ or } p \equiv 1 \mod 3. \end{cases}$$

$$\sharp D \cap \mathrm{Conj}(\sigma) = \begin{cases} p+3 & \text{if } p \equiv -1 \mod 4, \\ p+1 & \text{if } p \equiv 1 \mod 4, \end{cases}$$

$$\sharp D \cap \mathrm{Conj}(\tau) = \begin{cases} 2 & \text{if } p \geq 5 \text{ and } p \equiv -1 \mod 3, \\ 0 & \text{if } p = 3 \text{ or } p \equiv 1 \mod 3. \end{cases}$$

$$\sharp D \cap \mathrm{Conj}(u) = 0 \quad \text{if } p \geq 3.$$

4.

$$\sharp E\cap \mathrm{Conj}(\sigma)\leq \begin{cases} 30 & \textit{if } p\equiv \pm 1 \mod 5,\\ 18 & \textit{if } p\geq 5 \textit{ and } p\not\equiv \pm 1 \mod 5.\end{cases}$$

$$\sharp E\cap \mathrm{Conj}(\tau)\leq \begin{cases} 20 & \textit{if } p\equiv \pm 1 \mod 5,\\ 8 & \textit{if } p\geq 5 \textit{ and } p\not\equiv \pm 1 \mod 5.\end{cases}$$

$$\sharp E\cap \mathrm{Conj}(u)=0 \quad \textit{if } p\geq 5.$$

Proof. 1 is easy.

2. Note that if $p\geq 3$ then for any $x\in(\mathbb{Z}/p\mathbb{Z})^\times$ there exist $a,c\in\mathbb{Z}/p\mathbb{Z}$ such that $a^2+c^2=x^{-1}$. Thus we have $\begin{pmatrix} a & -xc\\ c & xa\end{pmatrix}\in\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ and $\begin{pmatrix} a & -xc\\ c & xa\end{pmatrix}^{-1}\begin{pmatrix} 0 & 1\\ -1 & 0\end{pmatrix}\begin{pmatrix} a & -xc\\ c & xa\end{pmatrix}=\begin{pmatrix} 0 & x\\ -x^{-1} & 0\end{pmatrix}$.

3. Assume $p\geq 3$. We show $\sharp\left\{\begin{pmatrix} x & y\\ -\lambda y & -x\end{pmatrix}\right\}\cap\mathrm{Conj}(\sigma)=p+1$. Put $D_1=\left\{\begin{pmatrix} x & y\\ \lambda y & x\end{pmatrix}\right\}\cap\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ and $D_2=\left\{\begin{pmatrix} x & y\\ -\lambda y & -x\end{pmatrix}\right\}\cap\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$. Since $p\geq 3$, we have $\mathrm{Conj}(\sigma)=\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})^{\mathrm{Tr}=0}$. Thus $\left\{\begin{pmatrix} x & y\\ -\lambda y & -x\end{pmatrix}\right\}\cap\mathrm{Conj}(\sigma)=D_2=D\setminus D_1$. We have $D=\mathrm{Ker}\,(\det:\left\{\begin{pmatrix} x & y\\ \lambda y & x\end{pmatrix},\begin{pmatrix} x & y\\ -\lambda y & -x\end{pmatrix}\right\}\longrightarrow(\mathbb{Z}/p\mathbb{Z})^\times)$ and $D_1=\mathrm{Ker}\,(\det:\left\{\begin{pmatrix} x & y\\ \lambda y & x\end{pmatrix}\right\}\longrightarrow(\mathbb{Z}/p\mathbb{Z})^\times)$. Since $\det:\left\{\begin{pmatrix} x & y\\ \lambda y & x\end{pmatrix}\right\}\longrightarrow(\mathbb{Z}/p\mathbb{Z})^\times$ is surjective, we have $\sharp D_1=(p^2-1)/(p-1)=p+1$ and $\sharp D_1=2(p^2-1)/(p-1)=2(p+1)$. Therefore $\sharp\left\{\begin{pmatrix} x & y\\ -\lambda y & -x\end{pmatrix}\right\}\cap\mathrm{Conj}(\sigma)=2(p+1)-(p+1)=p+1$.

4. In A_5, there are 15 elements of order 2 and 20 elements of order 3. In S_4, there are 9 elements of order 2 and 8 elements of order 3.

□

Next we calculate the number of elements conjugate to σ,τ,u in $A_1\subseteq\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$.

Lemma 4.10. *In* $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$*, we have*

$$A_1\cap\mathrm{Conj}(\sigma)=\left\{\sigma,\begin{pmatrix} -1 & 2\\ -1 & 1\end{pmatrix},\begin{pmatrix} 1 & 1\\ 2 & -1\end{pmatrix}\right\},$$

$$A_1\cap\mathrm{Conj}(\tau)=\left\{\begin{pmatrix} 2 & -1\\ -1 & -1\end{pmatrix},\begin{pmatrix} -1 & 1\\ 1 & 2\end{pmatrix}\right\},$$

$$A_1\cap\mathrm{Conj}(u)=\emptyset,$$

$$A_1\cap\mathrm{Conj}(u^2)=\emptyset,$$

where $A_1 = \langle \sigma, \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \rangle$. *Hence we have*

$$\begin{aligned} \sharp A_1 \cap \mathrm{Conj}(\sigma) &= 3, \\ \sharp A_1 \cap \mathrm{Conj}(\tau) &= 2, \\ \sharp A_1 \cap \mathrm{Conj}(u) &= 0, \\ \sharp A_1 \cap \mathrm{Conj}(u^2) &= 0. \end{aligned}$$

Proof. It follows from Lemma 4.6. □

5. Calculation of conjugate elements in $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$

We calculate the number of elements conjugate to σ, τ, u in $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$.

Lemma 5.1. *Let $n \geq 1$ be an integer. In $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ we have*

$$\sharp\mathrm{Conj}(\sigma) = \begin{cases} (p-1)p^{2n-1} & \textit{if } p \equiv -1 \mod 4, \\ (p+1)p^{2n-1} & \textit{if } p \equiv 1 \mod 4, \\ 3 & \textit{if } p = 2 \textit{ and } n = 1, \\ 3 \cdot 2^{2n-3} & \textit{if } p = 2 \textit{ and } n \geq 2, \end{cases}$$

$$\sharp\mathrm{Conj}(\tau) = \begin{cases} (p-1)p^{2n-1} & \textit{if } p \equiv -1 \mod 3, \\ (p+1)p^{2n-1} & \textit{if } p \equiv 1 \mod 3, \\ 4 \cdot 3^{2n-2} & \textit{if } p = 3, \end{cases}$$

$$\sharp\mathrm{Conj}(u) = \begin{cases} \frac{1}{2}(p^2-1)p^{2n-2} & \textit{if } p \geq 3, \\ 3 & \textit{if } p = 2 \textit{ and } n = 1, \\ 6 & \textit{if } p = 2 \textit{ and } n = 2, \\ 3 \cdot 2^{2n-4} & \textit{if } p = 2 \textit{ and } n \geq 3. \end{cases}$$

Proof. As we have seen in the proof of Lemma 4.1, we have an equality $\sharp\mathrm{Conj}(\alpha) = \sharp\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})/\sharp\mathrm{Z}(\alpha)$ for any $\alpha \in \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$. As is well known, $\sharp\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) = (p+1)(p-1)p^{3n-2}$. Now we calculate $\sharp\mathrm{Z}(\alpha)$ for $\alpha = \sigma,\ \tau,\ u$. For $\alpha = \sigma$, we have

$$\mathrm{Z}(\sigma) = (\mathbb{Z}/p^n\mathbb{Z}[\sigma])^{\det=1} \cong \begin{cases} \mathrm{W}_n(\mathbb{F}_{p^2})^{\mathrm{Norm}=1} & \text{if } p \equiv -1 \mod 4, \\ (\mathbb{Z}/p^n\mathbb{Z} \times \mathbb{Z}/p^n\mathbb{Z})^{\mathrm{Norm}=1} & \text{if } p \equiv 1 \mod 4, \\ (\mathbb{Z}_2[\sqrt{-1}]/2^n\mathbb{Z}_2[\sqrt{-1}])^{\mathrm{Norm}=1} & \text{if } p = 2, \end{cases}$$

hence

$$\sharp\mathrm{Z}(\sigma) = \begin{cases} (p+1)p^{n-1} & \text{if } p \equiv -1 \mod 4, \\ (p-1)p^{n-1} & \text{if } p \equiv 1 \mod 4, \\ 2 & \text{if } p = 2 \text{ and } n = 1, \\ 2^{n+1} & \text{if } p = 2 \text{ and } n \geq 2. \end{cases}$$

For $\alpha = \tau$, we have

$$\mathrm{Z}(\tau) = (\mathbb{Z}/p^n\mathbb{Z}[\tau])^{\det=1} \cong \begin{cases} \mathrm{W}_n(\mathbb{F}_{p^2})^{\mathrm{Norm}=1} & \text{if } p \equiv -1 \mod 3, \\ (\mathbb{Z}/p^n\mathbb{Z} \times \mathbb{Z}/p^n\mathbb{Z})^{\mathrm{Norm}=1} & \text{if } p \equiv 1 \mod 3, \\ (\mathbb{Z}_3[\zeta_3]/3^n\mathbb{Z}_3[\zeta_3])^{\mathrm{Norm}=1} & \text{if } p = 3, \end{cases}$$

where ζ_3 is a primitive third root of unity. Hence

$$\sharp \mathrm{Z}(\tau)=\begin{cases}(p+1)p^{n-1} & \text{if } p\equiv -1 \mod 3,\\ (p-1)p^{n-1} & \text{if } p\equiv 1 \mod 3,\\ 2\cdot 3^n & \text{if } p=3.\end{cases}$$

For $\alpha=u$, we have $\mathrm{Z}(u)=\left\{\begin{pmatrix}a & b\\ 0 & a\end{pmatrix} | a^2=1\right\}$, hence

$$\sharp \mathrm{Z}(u)=\begin{cases}2p^n & \text{if } p\geq 3,\\ 2 & \text{if } p=2 \text{ and } n=1,\\ 8 & \text{if } p=2 \text{ and } n=2,\\ 2^{n+2} & \text{if } p=2 \text{ and } n\geq 3,\end{cases}$$

as required.

□

Next we calculate the number of elements conjugate to u^{p^r} in $\mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$.

Lemma 5.2. *Assume $r\geq 0$ and $n\geq 1$. In $\mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$, we have*

$$\sharp \mathrm{Conj}(u^{p^r})=\begin{cases}\frac{1}{2}(p^2-1)p^{2n-2} & \textit{if } p\geq 3,\\ 3 & \textit{if } p=2 \textit{ and } n=1,\\ 6 & \textit{if } p=2 \textit{ and } n=2,\\ 3\cdot 2^{2n-4} & \textit{if } p=2 \textit{ and } n\geq 3.\end{cases}$$

Proof. We have $\mathrm{Z}(u^{p^r})=\left\{\begin{pmatrix}a & b\\ c & d\end{pmatrix} | p^r c=0,\ p^r(a-d)=0,\ ad-bc=1\right\}$, which maps surjectively mod p^n onto $\left\{\begin{pmatrix}a & b\\ 0 & a\end{pmatrix}\in \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})\right\}$, and each fiber of the mod p^n map consists of p^{3r} elements.

□

We study the fiber of the mod p^m map

$$\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})\cap \mathrm{Conj}(\alpha)\longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z})\cap \mathrm{Conj}(\alpha),$$

where $\alpha=\sigma,\tau,u^{p^r}$.

Take two integers m,n with $1\leq m\leq n$. As in Section 1, let

$$f_{n,m}:\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})\longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z})$$

be the mod p^m map. For $\alpha=\sigma,\tau,u^{p^r}$ $(r\geq 0)$, put

$$V_\alpha^{r+n,r+m}=\alpha^{-1}(f_{r+n,r+m}^{-1}(\alpha)\cap \mathrm{Conj}(\alpha))\subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z}).$$

When $\alpha=\sigma,\tau$, we always take $r=0$. We sometimes omit the superscripts r,n,m and simply write V_α.

Lemma 5.3. *Let $r\geq 0$ and $1\leq m<n$ be integers. Assume $n\leq 2m$. If $p\geq 3$, then each $V_\alpha^{r+n,r+m}$ ($\alpha=\sigma,\tau,\ u^{p^r}$) is a subgroup of $(1+p^{r+m}\mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z}))^{\det=1}\cong \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})^{\mathrm{Tr}=0}$ and isomorphic to $(\mathbb{Z}/p^{n-m}\mathbb{Z})^2$. If $p=2$, then $V_\sigma^{n,m}$ for $m\geq 2$ (resp. $V_\tau^{n,m}$ for any m, resp. $V_{u^{p^r}}^{r+n,r+m}$ for $m\geq 3$) is a subgroup of $(1+p^{r+m}\mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z}))^{\det=1}\cong \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})^{\mathrm{Tr}=0}$ and isomorphic to $(\mathbb{Z}/p^{n-m}\mathbb{Z})^2$. Explicitly:*

If $p \geq 3$ *or* ($p = 2$ *and* $m \geq 2$)*, we have*

$$V_{\sigma}^{n,m} = \left\{ 1 + p^m \begin{pmatrix} a & b \\ b & -a \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \right\} \cong \left\{ \begin{pmatrix} a & b \\ b & -a \end{pmatrix} \in \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \right\};$$

For any p *and* m*, we have*

$$V_{\tau}^{n,m} = \left\{ 1 + p^m \begin{pmatrix} a & b \\ b-a & -a \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \right\} \cong \left\{ \begin{pmatrix} a & b \\ b-a & -a \end{pmatrix} \in \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \right\};$$

If $p \geq 3$ *or* ($p = 2$ *and* $m \geq 3$)*, we have*

$$V_{u^{p^r}}^{r+n,r+m} = \left\{ 1 + p^{r+m} \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z}) \right\} \cong \left\{ \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} \in \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \right\}.$$

In particular, the inverse image of one element by the following maps consists of p^2 *elements:*

- mod $p^m : \mathrm{SL}_2(\mathbb{Z}/p^{m+1}\mathbb{Z}) \cap \mathrm{Conj}(\sigma) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z}) \cap \mathrm{Conj}(\sigma)$ *if* $p \geq 3$ *or* ($p = 2$ *and* $m \geq 2$)*,*
- mod $p^m : \mathrm{SL}_2(\mathbb{Z}/p^{m+1}\mathbb{Z}) \cap \mathrm{Conj}(\tau) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z}) \cap \mathrm{Conj}(\tau)$ *for any* p *and any* $m \geq 1$*,*
- mod $p^{r+m} : \mathrm{SL}_2(\mathbb{Z}/p^{r+m+1}\mathbb{Z}) \cap \mathrm{Conj}(u^{p^r}) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^{r+m}\mathbb{Z}) \cap \mathrm{Conj}(u^{p^r})$ *if* $p \geq 3$ *or* ($p = 2$ *and* $m \geq 3$)*.*

Proof. Assume $\alpha = \sigma$. We will show

$$f_{n,m}^{-1}(\sigma) \cap \mathrm{Conj}(\sigma) = \left\{ \begin{pmatrix} p^m a & 1 + p^m b \\ -1 + p^m b & -p^m a \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \right\}.$$

Looking at the determinant and the trace, we get one inclusion "$\subseteq$". In particular, we have $\sharp f_{n,m}^{-1}(\sigma') \cap \mathrm{Conj}(\sigma) \leq p^{2(n-m)}$ for any $\sigma' \in \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z}) \cap \mathrm{Conj}(\sigma)$. Since mod $p^m : \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \cap \mathrm{Conj}(\sigma) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z}) \cap \mathrm{Conj}(\sigma)$ is surjective, Lemma 5.1 shows that the other inclusion "$\supseteq$" holds if $p \geq 3$ or ($p = 2$ and $m \geq 2$).

We can show

$$f_{n,m}^{-1}(\tau) \cap \mathrm{Conj}(\tau) = \left\{ \begin{pmatrix} 1 + p^m a & 1 + p^m b \\ -1 + p^m(b-a) & -p^m a \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z}) \right\}$$

for any p, m and

$$f_{r+n,r+m}^{-1}(u^{p^r}) \cap \mathrm{Conj}(u^{p^r}) = \left\{ \begin{pmatrix} 1 + p^{r+m} a & p^r + p^{r+m} b \\ 0 & 1 - p^{r+m} a \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z}) \right\}$$

for $p \geq 3$ or ($p = 2$ and $n \geq 3$) similarly by using Lemma 5.1 and 5.2.

$\square$

Remark 5.4. *If* $p = 2$*, the inverse image of one element by the following maps consists of* 2 *elements:*

- mod $2 : \mathrm{SL}_2(\mathbb{Z}/2^2\mathbb{Z}) \cap \mathrm{Conj}(\sigma) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z}) \cap \mathrm{Conj}(\sigma)$,
- mod $2^{r+2} : \mathrm{SL}_2(\mathbb{Z}/2^{r+3}\mathbb{Z}) \cap \mathrm{Conj}(u^{2^r}) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/2^{r+2}\mathbb{Z}) \cap \mathrm{Conj}(u^{2^r})$,
- mod $2^{r+1} : \mathrm{SL}_2(\mathbb{Z}/2^{r+2}\mathbb{Z}) \cap \mathrm{Conj}(u^{2^r}) \longrightarrow \mathrm{SL}_2(\mathbb{Z}/2^{r+1}\mathbb{Z}) \cap \mathrm{Conj}(u^{2^r})$.

From now on, we always assume the hypothesis in Lemma 5.3 when we write $V_{\alpha}^{r+n,r+m}$ (for $\alpha = \sigma, \tau, u^{p^r}$ and their conjugates: defined below), so that $V_{\alpha}^{r+n,r+m}$ is a free $\mathbb{Z}/p^{n-m}\mathbb{Z}$-submodule of rank 2 of $(1+p^{r+m}\mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z}))^{\det=1} \cong \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})^{\mathrm{Tr}=0}$.

Lemma 5.5. *Let $r \geq 0$ and $1 \leq m < n$ be integers. Assume $n \leq 2m$. For $\alpha = \sigma,\, \tau,\, u^{p^r}$, we have*

$$V_\alpha^{r+n,r+m} = \{1 + p^m(X\alpha^{-1} - \alpha^{-1}X) | X \in \mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})\}.$$

Proof. If $X = \begin{pmatrix} x & y \\ z & w \end{pmatrix}$, we have

$$X\sigma^{-1} - \sigma^{-1}X = \begin{pmatrix} y+z & -x+w \\ w-x & -z-y \end{pmatrix},$$

$$X\tau^{-1} - \tau^{-1}X = \begin{pmatrix} y+z & -x+y+w \\ w-x-z & -z-y \end{pmatrix}$$

and

$$Xu^{-p^r} - u^{-p^r}X = \begin{pmatrix} p^r z & -p^r(x-w) \\ 0 & -p^r z \end{pmatrix}.$$

□

Lemma 5.6. *Let $r \geq 0$ and $1 \leq m < n$ be integers. For $\alpha = \sigma, \tau, u^{p^r}$, take an element $\alpha' \in \mathrm{Conj}(\alpha) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+m}\mathbb{Z})$. Suppose two elements $\alpha_1, \alpha_2 \in \mathrm{Conj}(\alpha) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ satisfy $\alpha_1 \equiv \alpha_2 \equiv \alpha' \mod p^{r+m}$. If $n \leq 2m$, then we have*

$${\alpha_1}^{-1}(f_{r+n,r+m}^{-1}(\alpha') \cap \mathrm{Conj}(\alpha)) = {\alpha_2}^{-1}(f_{r+n,r+m}^{-1}(\alpha') \cap \mathrm{Conj}(\alpha)).$$

Proof. It suffices to show the equality in the case $\alpha_2 = \alpha$. By the hypothesis, we have $\alpha_1 \in (f_{r+n,r+m}^{-1}(\alpha) \cap \mathrm{Conj}(\alpha)) = \alpha V_\alpha$. We see that ${\alpha_1}^{-1}(f_{r+n,r+m}^{-1}(\alpha) \cap \mathrm{Conj}(\alpha)) = {\alpha_1}^{-1}\alpha V_\alpha = \{{\alpha_1}^{-1}\alpha + (p^m {\alpha_1}^{-1}\alpha)(X\alpha^{-1} - \alpha^{-1}X)\} = \{{\alpha_1}^{-1}\alpha + p^m(X\alpha^{-1} - \alpha^{-1}X)\}$. Since $({\alpha_1}^{-1}\alpha)^{-1} = \alpha^{-1}\alpha_1 \in V_\alpha$, we have ${\alpha_1}^{-1}\alpha \in V_\alpha = \{1+p^m(X\alpha^{-1} - \alpha^{-1}X)\}$. Therefore $\alpha_1^{-1}(f_{r+n,r+m}^{-1}(\alpha) \cap \mathrm{Conj}(\alpha)) = \{1+p^m(X\alpha^{-1} - \alpha^{-1}X)\} = \alpha^{-1}(f_{r+n,r+m}^{-1}(\alpha) \cap \mathrm{Conj}(\alpha))$, as required.

□

In the above lemma, we define

$$V_{\alpha'}^{r+n,r+m} := {\alpha_1}^{-1}(f_{r+n,r+m}^{-1}(\alpha') \cap \mathrm{Conj}(\alpha)).$$

Note that we have $V_{\alpha'}^{r+n,r+m} = \{1 + p^m(X{\alpha'}^{-1} - {\alpha'}^{-1}X)\}$. For an element $g \in \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$, we have $V_{g^{-1}\alpha' g}^{r+n,r+m} = g^{-1}(V_{\alpha'}^{r+n,r+m})g$. We see that $V_{\alpha'}^{r+n,r+m}$ depends only on $\alpha' \mod p^{r+n-m}$. Thus we can define $V_{\alpha''}^{r+n,r+m}$ for $\alpha'' \in \mathrm{Conj}(\alpha) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n-m}\mathbb{Z})$.

For $\alpha = \sigma, \tau, u^{p^r}$ and their conjugates, we identify $V_\alpha^{r+n,r+m}$ with a free submodule of rank 2 of $\mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})$ using the isomorphisms in Lemma 5.3.

Lemma 5.7. *Let $1 \leq m < n$ be integers. Take an element $\alpha \in \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z})$ which is conjugate to σ or τ. Assume $n \leq 2m$. Then $V_\alpha^{n,m}$ is the orthogonal complement of $\mathbb{Z}/p^{n-m}\mathbb{Z}[\alpha]$ in $\mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})$ with respect to the pairing*

$$\mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \times \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \longrightarrow \mathbb{Z}/p^{n-m}\mathbb{Z} \;:\; (A, B) \mapsto \mathrm{Tr}\,(AB).$$

Proof. We have $V_\alpha^{n,m} = \{1 + p^m(X\alpha^{-1} - \alpha^{-1}X) \in 1 + p^m\mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z})\} \cong \{X\alpha^{-1} - \alpha^{-1}X \in \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})\}$ and it is orthogonal to $\mathbb{Z}/p^{n-m}\mathbb{Z}[\alpha]$. Since $\sharp V_\alpha^{n,m} = \sharp\mathbb{Z}/p^{n-m}\mathbb{Z}[\alpha] = p^{2(n-m)}$, they are the orthogonal complements of each other.

□

Note that

$$\mathbb{Z}/p^{n-m}\mathbb{Z}[\sigma] = \left\{ \begin{pmatrix} x & y \\ -y & x \end{pmatrix} \right\}$$

and

$$\mathbb{Z}/p^{n-m}\mathbb{Z}[\tau] = \left\{ \begin{pmatrix} x & y \\ -y & x-y \end{pmatrix} \right\}.$$

Lemma 5.8. *Let $r \geq 0$ and $1 \leq m < n$ be integers. Take an element $v = 1 + p^r \epsilon \in \mathrm{Conj}(u^{p^r}) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+m}\mathbb{Z})$. Assume $n \leq 2m$. Then $V_v^{r+n,r+m}$ is the orthogonal complement of $\mathbb{Z}/p^{n-m}\mathbb{Z}[\epsilon]$ in $\mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})$ with respect to the pairing*

$$\mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \times \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z}) \longrightarrow \mathbb{Z}/p^{n-m}\mathbb{Z} \ : \ (A, B) \mapsto \mathrm{Tr}\,(AB).$$

Proof. We have $V_v^{r+n,r+m} = \{1 + p^m(Xv^{-1} - v^{-1}X)\} = \{1 + p^m(X(1 - p^r\epsilon) - (1 - p^r\epsilon)X)\} = \{1 + p^{r+m}(X(-\epsilon) - (-\epsilon)X) \in 1 + p^{r+m}\mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})\} \cong \{X(-\epsilon) - (-\epsilon)X \in \mathrm{M}_2(\mathbb{Z}/p^{n-m}\mathbb{Z})\}$ and it is orthogonal to $\mathbb{Z}/p^{n-m}\mathbb{Z}[\epsilon]$. Since $\sharp V_v^{r+n,r+m} = \sharp \mathbb{Z}/p^{n-m}\mathbb{Z}[\epsilon] = p^{2(n-m)}$, they are the orthogonal complements of each other. □

Note that

$$\mathbb{Z}/p^{n-m}\mathbb{Z}[\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}] = \left\{ \begin{pmatrix} x & y \\ 0 & x \end{pmatrix} \right\}.$$

Next we study the condition for the equality $V_\alpha = V_{\alpha'}$, where $\alpha' \in \mathrm{Conj}(\alpha)$.

Lemma 5.9. *Let $n \geq 1$ be an integer. In $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$, we have*

$$\left\{ \begin{pmatrix} x & y \\ -y & x \end{pmatrix} \right\} \cap \mathrm{Conj}(\sigma)$$

$$= \begin{cases} \{\sigma, \sigma^{-1}\} \\ \quad \text{if } p \geq 3, \\ \{\sigma\} \\ \quad \text{if } p = 2 \text{ and } n = 1, \\ \left\{ \sigma, \begin{pmatrix} 2 & 1 \\ -1 & 2 \end{pmatrix} \right\} \\ \quad \text{if } p = 2 \text{ and } n = 2, \\ \left\{ \sigma, \begin{pmatrix} 0 & 1 + 2^{n-1} \\ -1 + 2^{n-1} & 0 \end{pmatrix}, \begin{pmatrix} 2^{n-1} & 1 \\ -1 & 2^{n-1} \end{pmatrix}, \begin{pmatrix} 2^{n-1} & 1 + 2^{n-1} \\ -1 + 2^{n-1} & 2^{n-1} \end{pmatrix} \right\} \\ \quad \text{if } p = 2 \text{ and } n \geq 3. \end{cases}$$

Corollary 5.10. *Let $1 \leq m < n$ be integers. Take two elements $\sigma', \sigma'' \in \mathrm{Conj}(\sigma) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z})$. Assume $p \geq 3$ and $n \leq 2m$. Then $V_{\sigma'}^{n,m} = V_{\sigma''}^{n,m}$ holds if and only if $\sigma' \equiv {\sigma''}^{\pm 1} \bmod p^{n-m}$.*

Proof. Replacing σ'' by its conjugate, we may assume $\sigma'' = \sigma$. By Lemma 5.7, we see that $\sigma' \bmod p^{n-m}$ is orthogonal to $V_{\sigma'} = V_\sigma$, and thus we have $\sigma' \bmod p^{n-m} \in \left\{ \begin{pmatrix} x & y \\ -y & x \end{pmatrix} \right\}$. Applying the previous lemma, we get the result. □

Corollary 5.11. *Let $2 \leq m < n$ be integers. Fix an element $\sigma' \in \mathrm{Conj}(\sigma) \subseteq \mathrm{SL}_2(\mathbb{Z}/2^m\mathbb{Z})$. Assume $n \leq 2m$. Then the number of the elements $\sigma'' \in \mathrm{Conj}(\sigma) \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{n-m}\mathbb{Z})$ satisfying $V^{n,m}_{\sigma''} = V^{n,m}_{\sigma'}$ is*

$$\begin{cases} 1 & \text{if } n-m=1, \\ 2 & \text{if } n-m=2, \\ 4 & \text{if } n-m \geq 3. \end{cases}$$

Lemma 5.12. *Let $n \geq 1$ be an integer. In $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$, we have*

$$\left\{ \begin{pmatrix} x & y \\ -y & x-y \end{pmatrix} \right\} \cap \mathrm{Conj}(\tau)$$

$$= \begin{cases} \{\tau, \tau^{-1}\} & \text{if } p \neq 3, \\ \{\tau\} & \text{if } p = 3 \text{ and } n = 1, \\ \left\{ \tau, \begin{pmatrix} 1+3^{n-1} & 1-3^{n-1} \\ -1+3^{n-1} & -3^{n-1} \end{pmatrix}, \begin{pmatrix} 1-3^{n-1} & 1+3^{n-1} \\ -1-3^{n-1} & 3^{n-1} \end{pmatrix} \right\} & \text{if } p = 3 \text{ and } n \geq 2. \end{cases}$$

Corollary 5.13. *Let $1 \leq m < n$ be integers. Take two elements $\tau', \tau'' \in \mathrm{Conj}(\tau) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^m\mathbb{Z})$. Assume $p \neq 3$ and $n \leq 2m$. Suppose $m \geq 2$ if $p = 2$. Then $V^{n,m}_{\tau'} = V^{n,m}_{\tau''}$ holds if and only if $\tau' \equiv \tau''^{\pm 1} \bmod p^{n-m}$.*

Corollary 5.14. *Let $1 \leq m < n$ be integers. Fix an element $\tau' \in \mathrm{Conj}(\tau) \subseteq \mathrm{SL}_2(\mathbb{Z}/3^m\mathbb{Z})$. Assume $n \leq 2m$. Then the number of the elements $\tau'' \in \mathrm{Conj}(\tau) \subseteq \mathrm{SL}_2(\mathbb{Z}/3^{n-m}\mathbb{Z})$ satisfying $V^{n,m}_{\tau''} = V^{n,m}_{\tau'}$ is*

$$\begin{cases} 1 & \text{if } n-m=1, \\ 3 & \text{if } n-m \geq 2. \end{cases}$$

Lemma 5.15. *Let $r \geq 0$ and $n \geq 1$ be integers. In $\mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$, we have*

$$\left\{ 1 + p^r \begin{pmatrix} x & y \\ 0 & x \end{pmatrix} \right\} \cap \mathrm{Conj}(u^{p^r})$$

$$= \begin{cases} \{u^{p^r s^2} | (p,s)=1\} & \text{if } p \geq 3, \\ \{u^{2^r}\} & \text{if } p = 2 \text{ and } n = 1, \\ \left\{ u^{2^r}, \begin{pmatrix} -1 & 2^r \\ 0 & -1 \end{pmatrix} \right\} & \text{if } p = 2 \text{ and } n = 2, \\ \left\{ u^{2^r s^2}, \begin{pmatrix} 1+2^{r+n-1} & 2^r s^2 \\ 0 & 1+2^{r+n-1} \end{pmatrix} | (2,s)=1 \right\} & \text{if } p = 2 \text{ and } n \geq 3. \end{cases}$$

We have $\sharp\{u^{p^r s^2} | (p,s)=1\} = \frac{1}{2}(p-1)p^{n-1}$ if $p \geq 3$, and $\sharp\left\{ u^{2^r s^2}, \begin{pmatrix} 1+2^{r+n-1} & 2^r s^2 \\ 0 & 1+2^{r+n-1} \end{pmatrix} | (2,s)=1 \right\} = 2^{n-2}$ *if $n \geq 3$.*

Corollary 5.16. *Let $r \geq 0$ and $1 \leq m < n$ be integers. Take two elements $v, v' \in \mathrm{Conj}(u^{p^r}) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+m}\mathbb{Z})$. Assume $p \geq 3$ and $n \leq 2m$. Then $V^{r+n,r+m}_v = V^{r+n,r+m}_{v'}$ holds if and only if $v \equiv v'^{s^2} \bmod p^{r+n-m}$ where $(p,s)=1$.*

Proof. Replacing v' by its conjugate, we may assume $v' = u^{p^r}$. Write $v = 1 + p^r\epsilon$. By Lemma 5.8, we see that $\epsilon \bmod p^{n-m}$ is orthogonal to $V_v = V_{u^{p^r}}$, and thus we have $\epsilon \bmod p^{n-m} \in \left\{\begin{pmatrix} x & y \\ 0 & x \end{pmatrix}\right\}$. Hence we have $v \bmod p^{r+n-m} \in \left\{1 + p^r \begin{pmatrix} x & y \\ 0 & x \end{pmatrix}\right\}$. Applying the previous lemma, we get the result.

□

Corollary 5.17. *Let $r \geq 0$ and $3 \leq m < n$ be integers. Fix an element $v \in \mathrm{Conj}(u^{2^r}) \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{r+m}\mathbb{Z})$. Assume $n \leq 2m$. Then the number of the elements $v' \in \mathrm{Conj}(u^{2^r}) \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{r+n-m}\mathbb{Z})$ satisfying $V_{v'}^{r+n,r+m} = V_v^{r+n,r+m}$ is*

$$\begin{cases} 1 & \text{if } n-m = 1, \\ 2 & \text{if } n-m = 2, \\ 2^{n-m-2} & \text{if } n-m \geq 3. \end{cases}$$

6. Control of inverse images

We control the number of elements conjugate to σ, τ, u^{p^r} contained in a slim subgroup H.

Let $n \geq 1$ be an integer and let H be a subgroup of $\mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$. For an integer $1 \leq s \leq n$, put

$$H_s := H \cap (1 + p^s\mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z})) = \mathrm{Ker}\,(\mathrm{mod} p^s : H \longrightarrow \mathrm{SL}_2(\mathbb{Z}/p^s\mathbb{Z})).$$

We identify H/H_s with $H \bmod p^s$. For two integers s, t with $1 \leq s \leq t \leq n$ and for $\alpha = \sigma, \tau, u^{p^r}$, let

$$f_{t,s}^{H,\alpha} : (H/H_t) \cap \mathrm{Conj}(\alpha) \longrightarrow (H/H_s) \cap \mathrm{Conj}(\alpha)$$

be the mod p^s map. Here we assume $s > r$ when $\alpha = u^{p^r}$.

Recall that a subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ is called a slim subgroup if $H \not\supseteq (1 + p^{n-1}\mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z}))^{\det=1}$, equivalently $\sharp H_{n-1} \leq p^2$. We prepare for controlling $\sharp H \cap \mathrm{Conj}(\alpha)$ for a slim subgroup H.

Lemma 6.1. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ be a slim subgroup. Take two integers s, t with $1 \leq t < s \leq n$. Assume $t \geq 2$ if $p = 2$. Then we have $\sharp H_t/H_s \leq p^{2(s-t)}$.*

Proof. Put $G = \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$. Take an integer i with $2 \leq i \leq n-1$. Assume $i \geq 3$ if $p = 2$. Then the p-th power map $\eta : G_{i-1}/G_i \longrightarrow G_i/G_{i+1}$ is surjective. As $\sharp G_{i-1}/G_i = \sharp G_i/G_{i+1} = p^3$, we see that η is an isomorphism. Let $\eta' : H_{i-1}/H_i \longrightarrow H_i/H_{i+1}$ be the p-th power map. The commutative diagram

$$\begin{tikzcd} H_{i-1}/H_i \arrow[r, "\subseteq"] \arrow[d, "\eta'"'] & G_{i-1}/G_i \arrow[d, "\eta"'] \\ H_i/H_{i+1} \arrow[r, "\subseteq"] & G_i/G_{i+1} \end{tikzcd}$$

shows that η' is injective. By the hypothesis $H \not\supseteq (1 + p^{n-1}\mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z}))^{\det=1}$, we have $H_{n-1}/H_n \subsetneq G_{n-1}/G_n$. Hence $\sharp H_{n-1}/H_n \leq p^2$. Therefore $\sharp H_1/H_2 \leq \sharp H_2/H_3 \leq \cdots \leq \sharp H_{n-1}/H_n \leq p^2$ if $p \geq 3$, while $\sharp H_2/H_3 \leq \sharp H_3/H_4 \leq \cdots \leq \sharp H_{n-1}/H_n \leq p^2$ if $p = 2$. Consequently, we get $\sharp H_t/H_s = \prod_{i=t+1}^{s} \sharp H_{i-1}/H_i \leq p^{2(s-t)}$.

□

From now to the end of this section, we use the letter α to denote any of σ, τ, u^{p^r}, where $r \geq 0$ be an integer. As usual, assume $r = 0$ if $\alpha = \sigma, \tau$.

Corollary 6.2. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a slim subgroup. Take two integers s, t with $1 \leq t < s \leq n$. Take an element $\alpha' \in \mathrm{Conj}(\alpha) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+t}\mathbb{Z})$. Assume $s \leq 2t$. When $p = 2$, further assume $r + t \geq 2$. If $H/H_{r+s} \supseteq V_{\alpha'}^{r+s,r+t}$, then $H_{r+t}/H_{r+s} = V_{\alpha'}^{r+s,r+t}$.*

Proof. If $H/H_{r+s} \supseteq V_{\alpha'}^{r+s,r+t}$, then $H_t/H_{r+s} \supseteq V_{\alpha'}^{r+s,r+t}$. By Lemma 5.3, we have $\sharp V_{\alpha'}^{r+s,r+t} = p^{2(s-t)}$. We have seen $\sharp H_{r+t}/H_{r+s} \leq p^{2(s-t)}$ in the previous lemma. Thus we get $H_{r+t}/H_{r+s} = V_{\alpha'}^{r+s,r+t}$.

□

Remark 6.3. *By using the diagram in the proof of Lemma 6.1, we can show Lemma 2.2.*

Lemma 6.4. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a slim subgroup. Take three integers s, t, i satisfying $1 \leq t < s \leq n$ and $i \geq 1$. Take an element $\alpha' \in \mathrm{Conj}(\alpha) \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+t}\mathbb{Z})$. Assume $s \leq 2t$ and $s + i \leq n$. When $p = 2$ and $\alpha = \tau$, further assume $s \leq 2t - 1$. If $H_{r+t}/H_{r+s} = V_{\alpha'}^{r+s,r+t}$, then $H_{r+t+i}/H_{r+s+i} = V_{\alpha'}^{r+s+i,r+t+i}$.*

Proof. Since $H_{r+t} \bmod p^{r+s} = \{1 + p^t(X\alpha'^{-1} - \alpha'^{-1}X) | X \in \mathrm{M}_2(\mathbb{Z}/p^{r+s}\mathbb{Z})\}$, we have $H \ni 1 + p^t(X\alpha'^{-1} - \alpha'^{-1}X) + p^{r+s}Y$ for some $Y \in \mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$.

Case $\alpha = \sigma$ or τ. In this case $r = 0$. Suppose $p \geq 3$ or ($p = 2$ and $s \leq 2t - 1$). Then we have $H \ni (1 + p^t(X\alpha'^{-1} - \alpha'^{-1}X) + p^sY)^{p^i} = 1 + p^{t+i}(X\alpha'^{-1} - \alpha'^{-1}X) + p^{s+i}Y'$ for some $Y' \in \mathrm{M}_2(\mathbb{Z}/p^n\mathbb{Z})$. Thus $H_{t+i}/H_{s+i} \supseteq \{1 + p^{t+i}(X\alpha'^{-1} - \alpha'^{-1}X)\} = V_{\alpha'}^{s+i,t+i}$. By Corollary 6.2, we have the equality $H_{t+i}/H_{s+i} = V_{\alpha'}^{s+i,t+i}$. Next suppose $\alpha = \sigma$, $p = 2$ and $s = 2t$. Notice that we are assuming $t \geq 2$ in this case. Replacing α' by its conjugate, we may assume $\alpha' = \sigma$. Then we have $H \ni 1 + 2^t \begin{pmatrix} a & b \\ b & -a \end{pmatrix} + 2^{2t}Y$, where $X\sigma^{-1} - \sigma^{-1}X = \begin{pmatrix} a & b \\ b & -a \end{pmatrix}$. Taking the second power, we have $H \ni 1 + 2^{t+1} \begin{pmatrix} a + 2^{t-1}(a^2 + b^2) & b \\ b & -a + 2^{t-1}(a^2 + b^2) \end{pmatrix} + 2^{2t+1}Y'$ for some $Y' \in \mathrm{M}_2(\mathbb{Z}/2^n\mathbb{Z})$. Put $R(a, b) := \begin{pmatrix} a + 2^{t-1}(a^2 + b^2) & b \\ b & -a + 2^{t-1}(a^2 + b^2) \end{pmatrix}$. The elements $R(a, b)$ generate $V_\sigma^{2t+1,t+1} \cong \left\{ \begin{pmatrix} x & y \\ y & -x \end{pmatrix} \in \mathrm{M}_2(\mathbb{Z}/2^t\mathbb{Z}) \right\}$ because $(1 + 2^{t-1})R(1, 0) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $R(0, 1) + 2^{t-1}R(1, 0) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Hence $H_{t+1}/H_{2t+1} \supseteq V_\sigma^{2t+1,t+1}$, and this implies $H_{t+1}/H_{2t+1} = V_\sigma^{2t+1,t+1}$ by Corollary 6.2. Therefore we get $H_{t+i}/H_{2t+i} = V_\sigma^{2t+i,t+i}$ as we have just seen.

Case $\alpha = u^{p^r}$. We may assume $\alpha' = u^{p^r}$. Thus we have $H \ni 1 + p^{r+t} \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} + p^{r+s}Y$, where $Xu^{-p^r} - u^{-p^r}X = p^r \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix}$. Suppose $p \geq 3$ or ($p = 2$ and $r + s \leq 2(r + t) - 1$). Then we have $H \ni (1 + p^{r+t} \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} + p^{r+s}Y)^{p^i} = 1 + p^{r+t+i} \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} + p^{r+s+i}Y'$ for some $Y' \in \mathrm{M}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$. Hence we get

$H_{r+t+i}/H_{r+s+i} = V_{u^{p^r}}^{r+s+i,r+t+i}$. Next suppose $p = 2$ and $r + s = 2(r + t)$, so that $r = 0$ and $s = 2t$. Notice that we are assuming $t \geq 3$ in this case. We have $H \ni (1 + 2^t \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} + 2^{2t}Y)^2 = 1 + 2^{t+1} \begin{pmatrix} a + 2^{t-1}a^2 & b \\ 0 & -a + 2^{t-1}a^2 \end{pmatrix} + 2^{2t+1}Y'$ for some $Y' \in \mathrm{M}_2(\mathbb{Z}/2^n\mathbb{Z})$. Put $S(a, b) := \begin{pmatrix} a + 2^{t-1}a^2 & b \\ 0 & -a + 2^{t-1}a^2 \end{pmatrix}$. The elements $S(a, b)$ generate $V_u^{2t+1,t+1} \cong \left\{ \begin{pmatrix} x & y \\ 0 & -x \end{pmatrix} \in \mathrm{M}_2(\mathbb{Z}/2^t\mathbb{Z}) \right\}$ because $(1 + 2^{t-1})S(1, 0) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $S(0, 1) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Therefore $H_{t+1}/H_{2t+1} = V_u^{2t+1,t+1}$ and $H_{t+i}/H_{2t+i} = V_u^{2t+i,t+i}$.

□

Lemma 6.5. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a subgroup. Take two integers t, i with $t \geq 1$, $i \geq 1$ and $t + i \leq n$. Take an element $\alpha' \in (H/H_{r+t}) \cap \mathrm{Conj}(\alpha)$. Assume $i \leq t$. If $H_{r+t}/H_{r+t+i} \neq V_{\alpha'}^{r+t+i,r+t}$, then we have $\sharp f_{r+t+i,r+t+1}((f_{r+t+i,r+t}^{H,\alpha})^{-1}(\alpha')) \leq p$.*

Proof. We show the lemma only when $r = 0$. Put $X := f_{t+i,t+1}((f_{t+i,t}^{H,\alpha})^{-1}(\alpha'))$. Suppose $(f_{t+i,t}^{H,\alpha})^{-1}(\alpha') \neq \emptyset$. Take an element $\widetilde{\alpha} \in (f_{t+i,t}^{H,\alpha})^{-1}(\alpha')$. The natural surjection mod p^{t+1} : $(f_{t+i,t}^{H,\alpha})^{-1}(\alpha') \longrightarrow X$ induces the following commutative diagram :

$$\begin{tikzcd}
V_{\alpha'}^{t+i,t} \cap (H/H_{t+i}) \arrow[r, "\mathrm{mod} p^{t+1}"] \arrow[d, "\subsetneq"'] & \widetilde{\alpha}^{-1}X \arrow[d, "\subseteq"'] \\
V_{\alpha'}^{t+i,t} \arrow[r, "\mathrm{mod} p^{t+1}"] \arrow[d, "\cong"'] & V_{\alpha'}^{t+1,t} \arrow[d, "\cong"'] \\
(\mathbb{Z}/p^i\mathbb{Z})^{\oplus 2} \arrow[r, "\mathrm{mod} p"] & (\mathbb{Z}/p\mathbb{Z})^{\oplus 2}.
\end{tikzcd}$$

All the horizontal maps in the above diagram are surjective. Since $V_{\alpha'}^{t+i,t} \cap (H/H_{t+i}) = V_{\alpha'}^{t+i,t} \cap (H_t/H_{t+i})$ is a proper subgroup of $V_{\alpha'}^{t+i,t} \cong (\mathbb{Z}/p^i\mathbb{Z})^{\oplus 2}$, we have $\sharp V_{\alpha'}^{t+i,t} \cap (H/H_{t+i}) \leq p^{2i-1}$. Since no proper subgroup of $(\mathbb{Z}/p^i\mathbb{Z})^{\oplus 2}$ maps surjectively mod$p$ onto $(\mathbb{Z}/p\mathbb{Z})^{\oplus 2}$, we get $\sharp \widetilde{\alpha}^{-1}X \leq p$.

□

Corollary 6.6. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a slim subgroup. Take two integers $i \geq 1$, $\delta \geq 0$ with $i + \delta \leq n$. Take an element $\alpha' \in (H/H_{r+i+\delta}) \cap \mathrm{Conj}(\alpha)$. Assume $2i + \delta \leq n$. When $p = 2$ and $\alpha = \tau$, further assume $\delta \geq 1$. If $H_{r+n-i}/H_{r+n} \neq V_{\alpha'}^{r+n,r+n-i}$, then we have $\sharp (f_{r+n,r+i+\delta}^{H,\alpha})^{-1}(\alpha') \leq p^{n-1-\delta}$.*

Proof. We show the corollary only when $r = 0$. For an integer t with $i \leq t \leq n$, put $X_t := f_{n,t}((f_{n,i}^{H,\alpha})^{-1}(\alpha')) \subseteq (f_{t,i}^{H,\alpha})^{-1}(\alpha')$. For $i \leq t \leq s \leq n$, let $f_{X_s,X_t} : X_s \longrightarrow X_t$ be the modp^t reduction map. Note that f_{X_s,X_t} is surjective. For each element $\alpha'' \in X_{n-i+1}$, we have $\sharp f_{X_n,X_{n-i+1}}^{-1}(\alpha'') \leq p^{2(i-1)}$ by Lemma 5.3. Thus we have $\sharp X_n \leq p^{2(i-1)} \sharp X_{n-i+1}$. For $i + \delta \leq t \leq n - i$ and $\alpha'' \in X_t$, we show $\sharp f_{X_{t+1},X_t}^{-1}(\alpha'') \leq p$. We have $f_{X_{t+1},X_t}^{-1}(\alpha'') = f_{t+i,t+1}(f_{X_{t+i},X_t}^{-1}(\alpha''))$. By the assumption $H_{n-i}/H_n \neq$

$V_{\alpha'}^{n,n-i}$ and by Lemma 6.4, we have $H_t/H_{t+i} \neq V_{\alpha''}^{t+i,t}$. Since $f_{X_{t+i},X_t}^{-1}(\alpha'') \subseteq (f_{t+i,t}^{H,\alpha})^{-1}(\alpha'')$, we have the inequality $\sharp f_{t+i,t+1}(f_{X_{t+i},X_t}^{-1}(\alpha'')) \leq p$ by Lemma 6.5. Hence we have $\sharp f_{X_{t+1},X_t}^{-1}(\alpha'') \leq p$ for $i+\delta \leq t \leq n-i$ and $\alpha'' \in X_t$. Thus we get $\sharp X_{t+1} \leq p \sharp X_t$ for $i+\delta \leq t \leq n-i$. Consequently we have $\sharp (f_{n,i}^{H,\alpha})^{-1}(\alpha') = \sharp X_n \leq p^{2(i-1)} \cdot p^{n-2i+1-\delta} \sharp X_i \leq p^{n-1-\delta}$.

□

Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a slim subgroup. Put

$$l := \begin{cases} \frac{n}{2} & \text{if } n \text{ is even,} \\ \frac{n-1}{2} & \text{if } n \text{ is odd.} \end{cases}$$

Define a decreasing sequence

$$Y_0 \supseteq Y_1 \supseteq \cdots \supseteq Y_i \supseteq Y_{i+1} \supseteq \cdots \supseteq Y_l$$

by

$$Y_i := \begin{cases} H \cap \mathrm{Conj}(\alpha) & \text{if } i = 0, \\ \{\alpha' \in H \cap \mathrm{Conj}(\alpha) | H_{r+n-i} = V_{\alpha'}^{r+n,r+n-i}\} & \text{if } 1 \leq i \leq l. \end{cases}$$

When $p \geq 3$, we use Y_i for $0 \leq i \leq l$; when $p = 2$, we use Y_i only for

$$\begin{cases} i = 0, 1 & \text{if } \alpha = \sigma \text{ and } 3 \leq n \leq 5, \\ 0 \leq i \leq l & \text{if } \alpha = \sigma \text{ and } n \geq 6, \\ i = 0,\ 3 \leq i \leq l & \text{if } \alpha = u^{2^r} \text{ and } n \geq 6, \end{cases}$$

so that the hypothesis in Lemma 5.3 is satisfied.

Lemma 6.7. *For three integers i, i', i'' satisfying $0 \leq i' < i \leq i'' \leq l$, we have*

$$(Y_{i'} \setminus Y_i) \bmod p^{r+i''} = (Y_{i'} \bmod p^{r+i''}) \setminus (Y_i \bmod p^{r+i''}).$$

Proof. The inclusion "$\supseteq$" is trivial. Now we show the other inclusion "$\subseteq$". Take an element $\alpha' \bmod p^{r+i''} \in (Y_{i'} \setminus Y_i) \bmod p^{r+i''}$ where $\alpha' \in Y_{i'} \setminus Y_i$. Assume $\alpha' \bmod p^{r+i''} \in Y_i \bmod p^{r+i''}$. Then we find an element $\alpha'' \in Y_i$ such that $\alpha' \equiv \alpha'' \bmod p^{r+i''}$. Since $i \leq i''$, we have $V_{\alpha'}^{r+n,r+n-i} = V_{\alpha''}^{r+n,r+n-i} = H_i$. But this contradicts the assumption $\alpha' \notin Y_i$. Hence we get the inclusion "$\subseteq$".

□

Proposition 6.8. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a slim subgroup. Take an integer s satisfying $1 \leq s \leq l$. Then we have*

$$\begin{aligned} &\sharp H \cap \mathrm{Conj}(\alpha) \\ \leq &(p^{2(n-l)} - p^{n-1}) \sharp (Y_l \bmod p^{r+l}) + (p^2 - 1) p^{n-1} \sum_{i=s}^{l-1} \sharp (Y_i \bmod p^{r+i}) \\ &+ p^{n-1} \sharp (H/H_{r+s}) \cap \mathrm{Conj}(\alpha). \end{aligned}$$

Proof. For simplicity, we show the inequality only when $r = 0$. Decompose

$$\sharp Y_0 = \sharp Y_l + \sum_{i=s+1}^{l} \sharp (Y_{i-1} \setminus Y_i) + \sharp (Y_0 \setminus Y_s).$$

We have

$$\sharp Y_l \leq p^{2(n-l)} \sharp (Y_l \bmod p^l)$$

by Lemma 5.3. We see

$$\sharp(Y_{i-1}\setminus Y_i) \le p^{n-1}\sharp((Y_{i-1}\setminus Y_i) \bmod p^i) = p^{n-1}(\sharp(Y_{i-1} \bmod p^i) - \sharp(Y_i \bmod p^i))$$

and

$$\sharp(Y_0\setminus Y_s) \le p^{n-1}\sharp((Y_0\setminus Y_s) \bmod p^s) = p^{n-1}(\sharp(Y_0 \bmod p^s) - \sharp(Y_s \bmod p^s))$$

by the definition of Y_i, Corollary 6.6 and Lemma 6.7. Hence

$$\begin{aligned}\sharp Y_0 \le& p^{2(n-l)}\sharp(Y_l \bmod p^l) + p^{n-1}\sum_{i=s+1}^{l}(\sharp(Y_{i-1} \bmod p^i) - \sharp(Y_i \bmod p^i))\\ &+ p^{n-1}(\sharp(Y_0 \bmod p^s) - \sharp(Y_s \bmod p^s))\\ =&(p^{2(n-l)} - p^{n-1})\sharp(Y_l \bmod p^l) + p^{n-1}\sum_{i=s}^{l-1}(\sharp(Y_i \bmod p^{i+1}) - \sharp(Y_i \bmod p^i))\\ &+ p^{n-1}\sharp(Y_0 \bmod p^s).\end{aligned}$$

By Lemma 5.3, we see

$$\sharp(Y_i \bmod p^{i+1}) - \sharp(Y_i \bmod p^i) \le (p^2-1)\sharp(Y_i \bmod p^i).$$

Since

$$Y_0 \bmod p^s = (H/H_s) \cap \mathrm{Conj}(\alpha),$$

we get the desired inequality.

□

We control the number of elements conjugate to σ in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ when $p \ge 3$.

For $p \ge 3$, define a sequence $\{a(\sigma,p)_n\}_{n\ge 2}$ as follows:

$$a(\sigma,p)_n := 2p^{2(n-l)} + 2(l-1)(p^2-1)p^{n-1},$$

where $n = 2l$ or $2l+1$.

Corollary 6.9. *Let $n \ge 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ be a slim subgroup. Assume $p \ge 3$. Then we have*

$$\sharp H \cap \mathrm{Conj}(\sigma) \le a(\sigma,p)_n + p^{n-1}(\sharp(H/H_1) \cap \mathrm{Conj}(\sigma) - 2).$$

Proof. By Corollary 5.10, we have

$$\sharp(Y_i \bmod p^i) \le 2$$

for $1 \le i \le l$. Applying Proposition 6.8 (put $s = 1$), we get

$$\begin{aligned}&\sharp H \cap \mathrm{Conj}(\sigma)\\ \le&(p^{2(n-l)} - p^{n-1})\cdot 2 + (p^2-1)p^{n-1}\cdot 2(l-1) + p^{n-1}\sharp(H/H_1)\cap \mathrm{Conj}(\sigma)\\ =&a(\sigma,p)_n + p^{n-1}(\sharp(H/H_1) \cap \mathrm{Conj}(\sigma) - 2).\end{aligned}$$

□

We control the number of elements conjugate to τ in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ when $p \ge 3$.

For $p \ge 5$, define a sequence $\{a(\tau,p)_n\}_{n\ge 2}$ by

$$a(\tau,p)_n := a(\sigma,p)_n.$$

Corollary 6.10. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$ be a slim subgroup. Assume $p \geq 5$. Then we have*

$$\sharp H \cap \mathrm{Conj}(\tau) \leq a(\tau,p)_n + p^{n-1}(\sharp(H/H_1) \cap \mathrm{Conj}(\tau) - 2).$$

Proof. The same as the proof of Corollary 6.9.

□

Define a sequence $\{a(\tau,3)_n\}_{n\geq 2}$ as follows:

$$a(\tau,3)_n := \begin{cases} 3^2 & \text{if } n = 2, \\ (4n-11)\cdot 3^n & \text{if } n = 2l \geq 4, \\ (4n-9)\cdot 3^n & \text{if } n = 2l+1. \end{cases}$$

Corollary 6.11. *Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/3^n\mathbb{Z})$ be a slim subgroup. Then we have*

$$\sharp H \cap \mathrm{Conj}(\tau) \leq a(\tau,3)_n + 3^{n-1}(\sharp(H/H_1) \cap \mathrm{Conj}(\tau) - 1).$$

Proof. By Corollary 5.14, we have

$$\sharp(Y_i \bmod p^i) \leq \begin{cases} 1 & \text{if } i = 1, \\ 3 & \text{if } i \geq 2. \end{cases}$$

Applying Proposition 6.8 (put $s = 1$), we get

$$\begin{aligned} &\sharp H \cap \mathrm{Conj}(\tau) \\ \leq &\begin{cases} (3^{2(n-l)} - 3^{n-1})\cdot 1 + 3^{n-1}\sharp(H/H_1) \cap \mathrm{Conj}(\tau) \\ \text{ if } l = 1, \\ (3^{2(n-l)} - 3^{n-1})\cdot 3 + (3^2-1)\cdot 3^{n-1}\cdot 3(l-2) + (3^2-1)\cdot 3^{n-1}\cdot 1 \\ +3^{n-1}\sharp(H/H_1) \cap \mathrm{Conj}(\tau) \\ \text{ if } l \geq 2 \end{cases} \\ = &a(\tau,3)_n + 3^{n-1}(\sharp(H/H_1) \cap \mathrm{Conj}(\tau) - 1). \end{aligned}$$

□

We control the number of elements conjugate to u^{p^r} in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$.

For $p \geq 3$, define a sequence $\{a(u,p)_n\}_{n\geq 2}$ as follows:

$$a(u,p)_n := \begin{cases} \frac{1}{2}(p-1)(2\cdot p^{3l-1} - p^n) & \text{if } n = 2l, \\ \frac{1}{2}(p-1)(p^{3l+1} + p^{3l} - p^n) & \text{if } n = 2l+1. \end{cases}$$

Corollary 6.12. *Let $r \geq 0$, $n \geq 2$ be integers and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p^{r+n}\mathbb{Z})$ be a slim subgroup. Assume $p \geq 3$. Then we have*

$$\sharp H \cap \mathrm{Conj}(u^{p^r}) \leq a(u,p)_n + p^{n-1}(\sharp(H/H_{r+1}) \cap \mathrm{Conj}(u^{p^r}) - \frac{1}{2}(p-1)).$$

Proof. By Corollary 5.16, we have

$$\sharp(Y_i \bmod p^{r+i}) \leq \frac{1}{2}(p-1)p^{i-1}$$

for $1 \leq i \leq l$. Applying Proposition 6.8 (put $s = 1$), we get

$$\begin{aligned}&\sharp H \cap \mathrm{Conj}(u^{p^r})\\ \leq&(p^{2(n-l)} - p^{n-1}) \cdot \frac{1}{2}(p-1)p^{l-1} + (p^2-1)p^{n-1}\sum_{i=1}^{l-1}\frac{1}{2}(p-1)p^{i-1}\\ &+ p^{n-1}\sharp(H/H_{r+1}) \cap \mathrm{Conj}(u^{p^r})\\ =&a(u,p)_n + p^{n-1}(\sharp(H/H_{r+1}) \cap \mathrm{Conj}(u^{p^r}) - \frac{1}{2}(p-1)).\end{aligned}$$

□

Define a sequence $\{a(u,2)_n\}_{n\geq 6}$ as follows:

$$a(u,2)_n := \begin{cases} 2^{3l-1} - 2^{n+1} & \text{if } n = 2l, \\ 3 \cdot 2^{3l-1} - 2^{n+1} & \text{if } n = 2l+1. \end{cases}$$

Corollary 6.13. *Let $r \geq 0$, $n \geq 6$ be integers and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{r+n}\mathbb{Z})$ be a slim subgroup. Then we have*

$$\sharp H \cap \mathrm{Conj}(u^{2^r}) \leq a(u,2)_n + 2^{n-1}(\sharp(H/H_{r+3}) \cap \mathrm{Conj}(u^{2^r}) - 2).$$

Proof. By Corollary 5.17, we have

$$\sharp(Y_i \bmod 2^{r+i}) \leq 2^{i-2}$$

for $3 \leq i \leq l$. Applying Proposition 6.8 (put $s = 3$), we get

$$\begin{aligned}&\sharp H \cap \mathrm{Conj}(u^{2^r})\\ \leq&(2^{2(n-l)} - 2^{n-1}) \cdot 2^{l-2} + (2^2-1) \cdot 2^{n-1}\sum_{i=3}^{l-1} 2^{i-2} + 2^{n-1}\sharp(H/H_{r+3}) \cap \mathrm{Conj}(u^{2^r})\\ =&a(u,2)_n + 2^{n-1}(\sharp(H/H_{r+3}) \cap \mathrm{Conj}(u^{2^r}) - 2).\end{aligned}$$

□

We control the number of elements conjugate to σ in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^n\mathbb{Z})$.

Define a sequence $\{a(\sigma,2)_n\}_{n\geq 3}$ as follows:

$$a(\sigma,2)_n := \begin{cases} 2^3 & \text{if } n = 3, \\ 2^5 & \text{if } n = 4, \\ 3(l-2) \cdot 2^{n+1} & \text{if } n = 2l \geq 6, \\ (3l-4) \cdot 2^{n+1} & \text{if } n = 2l+1 \geq 5. \end{cases}$$

Proposition 6.14. *Let $n \geq 3$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^n\mathbb{Z})$ be a slim subgroup. Then we have*

$$\sharp H \cap \mathrm{Conj}(\sigma) \leq a(\sigma,2)_n + 2^{n-2}(\sharp(H/H_2) \cap \mathrm{Conj}(\sigma) - 2).$$

Proof. Decompose

$$\sharp Y_0 = \sharp Y_1 + \sharp(Y_0 \setminus Y_1).$$

We have

$$\sharp Y_1 \leq 2^{2(n-2)}\sharp(Y_1 \bmod 2^2)$$

by Lemma 5.3. We see

$$\sharp(Y_0 \setminus Y_1) \leq 2^{n-2}\sharp((Y_0 \setminus Y_1) \bmod 2^2) = 2^{n-2}(\sharp(Y_0 \bmod 2^2) - \sharp(Y_1 \bmod 2^2))$$

by the definition of Y_1, Corollary 6.6 and Lemma 6.7. Hence

$$\sharp Y_0 \leq (2^{2(n-2)} - 2^{n-2})\sharp(Y_1 \bmod 2^2) + 2^{n-2}\sharp(Y_0 \bmod 2^2).$$

As

$$\sharp(Y_1 \bmod 2^2) \leq 2$$

by Corollary 5.11 and Remark 5.4, we get

$$\sharp Y_0 \leq (2^{2(n-2)} - 2^{n-2})\cdot 2 + 2^{n-2}\sharp(Y_0 \bmod 2^2) = 2^{2n-3} + 2^{n-2}(\sharp(H/H_2) \cap \mathrm{Conj}(\sigma) - 2).$$

This shows the desired inequality for $n = 3, 4, 5$.

Assume $n \geq 6$, so that we have $l \geq 3$. Decompose

$$\sharp Y_0 = \sharp Y_l + \sum_{i=1}^{l} \sharp(Y_{i-1} \setminus Y_i).$$

We have

$$\sharp Y_l \leq 2^{2(n-l)}\sharp(Y_l \bmod 2^l)$$

by Lemma 5.3. We see

$$\begin{aligned}&\sharp(Y_{i-1} \setminus Y_i)\\ \leq &\begin{cases} 2^{n-1}\sharp((Y_{i-1} \setminus Y_i) \bmod 2^i) = 2^{n-1}(\sharp(Y_{i-1} \bmod 2^i) - \sharp(Y_i \bmod 2^i)) & \text{if } i \geq 2, \\ 2^{n-2}\sharp((Y_0 \setminus Y_1) \bmod 2^2) = 2^{n-2}(\sharp(Y_0 \bmod 2^2) - \sharp(Y_1 \bmod 2^2)) & \text{if } i = 1 \end{cases}\end{aligned}$$

by the definition of Y_i, Corollary 6.6 and Lemma 6.7. Hence

$$\begin{aligned}\sharp Y_0 \leq &2^{2(n-l)}\sharp(Y_l \bmod 2^l) + 2^{n-1}\sum_{i=2}^{l}(\sharp(Y_{i-1} \bmod 2^i) - \sharp(Y_i \bmod 2^i))\\ &+ 2^{n-2}(\sharp(Y_0 \bmod 2^2) - \sharp(Y_1 \bmod 2^2))\\ =&(2^{2(n-l)} - 2^{n-1})\sharp(Y_l \bmod 2^l) + 2^{n-1}\sum_{i=2}^{l-1}(\sharp(Y_i \bmod 2^{i+1}) - \sharp(Y_i \bmod 2^i))\\ &+ 2^{n-2}\sharp(Y_1 \bmod 2^2) + 2^{n-2}\sharp(Y_0 \bmod 2^2).\end{aligned}$$

By Lemma 5.3, we see

$$\sharp(Y_i \bmod 2^{i+1}) - \sharp(Y_i \bmod 2^i) \leq (2^2 - 1)\sharp(Y_i \bmod 2^i) = 3\sharp(Y_i \bmod 2^i)$$

for $2 \leq i \leq l-1$. Thus we have

$$\begin{aligned}\sharp Y_0 \leq &(2^{2(n-l)} - 2^{n-1})\sharp(Y_l \bmod 2^l) + 3 \cdot 2^{n-1}(\sum_{i=2}^{l-1}\sharp(Y_i \bmod 2^i))\\ &+ 2^{n-2}\sharp(Y_1 \bmod 2^2) + 2^{n-2}\sharp(Y_0 \bmod 2^2).\end{aligned}$$

As

$$\sharp(Y_i \bmod 2^i) \leq \begin{cases} 2 & \text{if } i = 2, \\ 4 & \text{if } i \geq 3 \end{cases}$$

and

$$\sharp(Y_1 \bmod 2^2) \leq 2$$

by Corollary 5.11 and Remark 5.4, we get

$$\begin{aligned}\sharp Y_0 \leq & (2^{2(n-l)} - 2^{n-1}) \cdot 4 + 3 \cdot 2^{n-1} \cdot 4(l-3) + 3 \cdot 2^{n-1} \cdot 2 + 2^{n-2} \cdot 2 \\ & + 2^{n-2} \sharp (Y_0 \bmod 2^2) \\ = & a(\sigma, 2)_n + 2^{n-2}(\sharp(H/H_2) \cap \mathrm{Conj}(\sigma) - 2).\end{aligned}$$

□

We also use a slightly different way to control $\sharp H \cap \mathrm{Conj}(\tau)$ and $\sharp H \cap \mathrm{Conj}(u^{p^r})$ when $p = 2$.

Let $n \geq 2$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{r+n}\mathbb{Z})$ be a slim subgroup. Put

$$l' := \begin{cases} \frac{n}{2} & \text{if } n \text{ is even}, \\ \frac{n+1}{2} & \text{if } n \text{ is odd}. \end{cases}$$

Define a decreasing sequence

$$Z_0 \supseteq Z_1 \supseteq \cdots \supseteq Z_i \supseteq Z_{i+1} \supseteq \cdots \supseteq Z_{l'-1}$$

by

$$Z_i := \begin{cases} H \cap \mathrm{Conj}(\alpha) & \text{if } i = 0, \\ \{\alpha' \in H \cap \mathrm{Conj}(\alpha) | H_{r+n-i} = V_{\alpha'}^{r+n, r+n-i}\} & \text{if } 1 \leq i \leq l' - 1. \end{cases}$$

When $p = 2$, we use Z_i only for

$$\begin{cases} i = 0,\ 2 \leq i \leq l' - 1 & \text{if } \alpha = \tau \text{ and } n \geq 5, \\ i = 0, 1 & \text{if } \alpha = u^{2^r} \text{ and } 4 \leq n \leq 6, \\ i = 0, 1,\ 3 \leq i \leq l' - 1 & \text{if } \alpha = u^{2^r} \text{ and } n \geq 7, \end{cases}$$

so that the hypothesis in Lemma 5.3 is satisfied.

Lemma 6.15. *For three integers i, i', i'' satisfying $0 \leq i' < i \leq i'' \leq l$, we have*

$$(Z_{i'} \setminus Z_i) \bmod p^{r+i''} = (Z_{i'} \bmod p^{r+i''}) \setminus (Z_i \bmod p^{r+i''}).$$

Proof. The same as the proof of Lemma 6.7.

□

We control the number of elements conjugate to τ in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^n\mathbb{Z})$.

Define a sequence $\{a(\tau, 2)_n\}_{n \geq 5}$ as follows:

$$a(\tau, 2)_n := \begin{cases} (3l' - 5) \cdot 2^{n+1} & \text{if } n = 2l', \\ (3l' - 7) \cdot 2^{n+1} & \text{if } n = 2l' - 1. \end{cases}$$

Proposition 6.16. *Let $n \geq 5$ be an integer and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^n\mathbb{Z})$ be a slim subgroup. Then we have*

$$\sharp H \cap \mathrm{Conj}(\tau) \leq a(\tau, 2)_n + 2^{n-2}(\sharp(H/H_3) \cap \mathrm{Conj}(\tau) - 8).$$

Proof. Decompose

$$\sharp Z_0 = \sharp Z_{l'-1} + \sum_{i=3}^{l'-1} \sharp(Z_{i-1} \setminus Z_i) + \sharp(Z_0 \setminus Z_2).$$

We have

$$\sharp Z_{l'-1} \leq 2^{2(n-l')} \sharp(Z_{l'-1} \bmod 2^{l'})$$

by Lemma 5.3. By the definition of Z_i, Corollary 6.6 and Lemma 6.15, we see

$$\begin{aligned}\sharp(Z_{i-1}\setminus Z_i) \leq& 2^{n-2}\sharp((Z_{i-1}\setminus Z_i) \bmod 2^{i+1})\\ =&2^{n-2}(\sharp(Z_{i-1} \bmod 2^{i+1}) - \sharp(Z_i \bmod 2^{i+1}))\end{aligned}$$

for $3 \leq i \leq l'-1$ and

$$\sharp(Z_0\setminus Z_2) \leq 2^{n-2}\sharp((Z_0\setminus Z_2) \bmod 2^3) = 2^{n-2}(\sharp(Z_0 \bmod 2^3) - \sharp(Z_2 \bmod 2^3)).$$

Hence

$$\begin{aligned}\sharp Z_0 \leq& 2^{2(n-l')}\sharp(Z_{l'-1} \bmod 2^{l'}) + 2^{n-2}\sum_{i=3}^{l'-1}(\sharp(Z_{i-1} \bmod 2^{i+1}) - \sharp(Z_i \bmod 2^{i+1}))\\ &+ 2^{n-2}(\sharp(Z_0 \bmod 2^3) - \sharp(Z_2 \bmod 2^3))\\ =&(2^{2(n-l')} - 2^{n-2})\sharp(Z_{l'-1} \bmod 2^{l'})\\ &+ 2^{n-2}\sum_{i=2}^{l'-2}(\sharp(Z_i \bmod 2^{i+2}) - \sharp(Z_i \bmod 2^{i+1})) + 2^{n-2}\sharp(Z_0 \bmod 2^3).\end{aligned}$$

By Lemma 5.3, we see

$$\sharp(Z_i \bmod 2^{i+2}) - \sharp(Z_i \bmod 2^{i+1}) \leq (2^2-1)\sharp(Z_i \bmod 2^{i+1}) = 3\sharp(Z_i \bmod 2^{i+1})$$

for $2 \leq i \leq l'-2$. Therefore

$$\begin{aligned}\sharp Z_0 \leq&(2^{2(n-l')} - 2^{n-2})\sharp(Z_{l'-1} \bmod 2^{l'}) + 3\cdot 2^{n-2}\sum_{i=2}^{l'-2}\sharp(Z_i \bmod 2^{i+1})\\ &+ 2^{n-2}\sharp(Z_0 \bmod 2^3).\end{aligned}$$

By Corollary 5.13 and Lemma 5.3, we have

$$\sharp(Z_i \bmod 2^{i+1}) \leq 2^3$$

for $2 \leq i \leq l'-1$. Consequently, we get

$$\begin{aligned}\sharp Z_0 &\leq (2^{2(n-l')} - 2^{n-2})\cdot 2^3 + 3\cdot 2^{n-2}(l'-3)\cdot 2^3 + 2^{n-2}\sharp(Z_0 \bmod 2^3)\\ &= a(\tau,2)_n + 2^{n-2}(\sharp(H/H_3)\cap \mathrm{Conj}(\tau) - 8).\end{aligned}$$

□

We find another control of the number of elements conjugate to u^{2^r} in a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{r+n}\mathbb{Z})$. It is better than a previous one for a small n.

Define a sequence $\{b(u,2)_n\}_{n\geq 4}$ as follows:

$$b(u,2)_n := \begin{cases} 3\cdot 2^{3l'-2} - 2^{n+1} & \text{if } n = 2l',\\ 2^{3l'-2} - 2^{n+1} & \text{if } n = 2l'-1.\end{cases}$$

Proposition 6.17. *Let $r \geq 0$, $n \geq 4$ be integers and let $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{r+n}\mathbb{Z})$ be a slim subgroup. Then we have*

$$\sharp H \cap \mathrm{Conj}(u^{2^r}) \leq b(u,2)_n + 2^{n-3}(\sharp(H/H_{r+3})\cap \mathrm{Conj}(u^{2^r}) - 4).$$

Proof. Decompose

$$\sharp Z_0 = \sharp Z_1 + \sharp(Z_0\setminus Z_1).$$

We have

$$\sharp Z_1 \leq 2^{2(n-3)}\sharp(Z_1 \bmod 2^{r+3})$$

by Lemma 5.3. We see

$$\sharp(Z_0 \setminus Z_1) \leq 2^{n-3}\sharp((Z_0 \setminus Z_1) \bmod 2^{r+3}) = 2^{n-3}(\sharp(Z_0 \bmod 2^{r+3}) - \sharp(Z_1 \bmod 2^{r+3}))$$

by the definition of Z_1, Corollary 6.6 and Lemma 6.15. Hence

$$\sharp Z_0 \leq (2^{2(n-3)} - 2^{n-3})\sharp(Z_1 \bmod 2^{r+3}) + 2^{n-3}\sharp(Z_0 \bmod 2^{r+3}).$$

As

$$\sharp(Z_1 \bmod 2^{r+3}) \leq 2^2$$

by Corollary 5.17 and Remark 5.4, we get

$$\begin{aligned}\sharp Z_0 &\leq (2^{2(n-3)} - 2^{n-3}) \cdot 2^2 + 2^{n-3}\sharp(Z_0 \bmod 2^{r+3}) \\ &= 2^{2n-4} + 2^{n-3}(\sharp(H/H_{r+3}) \cap \mathrm{Conj}(u^{2^r}) - 4).\end{aligned}$$

This shows the desired inequality for $n = 4, 5, 6$.

Assume $n \geq 7$, so that we have $l' \geq 4$. Decompose

$$\sharp Z_0 = \sharp Z_{l'-1} + \sum_{i=4}^{l'-1} \sharp(Z_{i-1} \setminus Z_i) + \sharp(Z_1 \setminus Z_3) + \sharp(Z_0 \setminus Z_1).$$

We have

$$\sharp Z_{l'-1} \leq 2^{2(n-l')}\sharp(Z_{l'-1} \bmod 2^{r+l'})$$

by Lemma 5.3. By the definition of Z_i, Corollary 6.6 and Lemma 6.15, we see

$$\begin{aligned}\sharp(Z_{i-1} \setminus Z_i) \leq& 2^{n-2}\sharp((Z_{i-1} \setminus Z_i) \bmod 2^{r+i+1}) \\ =& 2^{n-2}(\sharp(Z_{i-1} \bmod 2^{r+i+1}) - \sharp(Z_i \bmod 2^{r+i+1}))\end{aligned}$$

for $4 \leq i \leq l' - 1$,

$$\sharp(Z_1 \setminus Z_3) \leq 2^{n-2}\sharp((Z_1 \setminus Z_3) \bmod 2^{r+4}) = 2^{n-2}(\sharp(Z_1 \bmod 2^{r+4}) - \sharp(Z_3 \bmod 2^{r+4}))$$

and

$$\sharp(Z_0 \setminus Z_1) \leq 2^{n-3}\sharp((Z_0 \setminus Z_1) \bmod 2^{r+3}) = 2^{n-3}(\sharp(Z_0 \bmod 2^{r+3}) - \sharp(Z_1 \bmod 2^{r+3})).$$

Hence

$$\begin{aligned}\sharp Z_0 \leq& 2^{2(n-l')}\sharp(Z_{l'-1} \bmod 2^{r+l'}) + 2^{n-2}\sum_{i=4}^{l'-1}(\sharp(Z_{i-1} \bmod 2^{r+i+1}) - \sharp(Z_i \bmod 2^{r+i+1})) \\ &+ 2^{n-2}(\sharp(Z_1 \bmod 2^{r+4}) - \sharp(Z_3 \bmod 2^{r+4})) \\ &+ 2^{n-3}(\sharp(Z_0 \bmod 2^{r+3}) - \sharp(Z_1 \bmod 2^{r+3})) \\ =& (2^{2(n-l')} - 2^{n-2})\sharp(Z_{l'-1} \bmod 2^{r+l'}) \\ &+ 2^{n-2}\sum_{i=3}^{l'-2}(\sharp(Z_i \bmod 2^{r+i+2}) - \sharp(Z_i \bmod 2^{r+i+1})) \\ &+ 2^{n-3}(2\sharp(Z_1 \bmod 2^{r+4}) - \sharp(Z_1 \bmod 2^{r+3})) + 2^{n-3}\sharp(Z_0 \bmod 2^{r+3}).\end{aligned}$$

By Lemma 5.3, we see

$$\begin{aligned}\sharp(Z_i \bmod 2^{r+i+2}) - \sharp(Z_i \bmod 2^{r+i+1}) \leq& (2^2 - 1)\sharp(Z_i \bmod 2^{r+i+1}) \\ =& 3\sharp(Z_i \bmod 2^{r+i+1})\end{aligned}$$

for $3 \leq i \leq l' - 2$ and

$$2\sharp(Z_1 \bmod 2^{r+4}) - \sharp(Z_1 \bmod 2^{r+3}) \leq (2 \cdot 2^2 - 1)\sharp(Z_1 \bmod 2^{r+3}) = 7\sharp(Z_1 \bmod 2^{r+3}).$$

Therefore

$$\begin{aligned}\sharp Z_0 \leq& (2^{2(n-l')} - 2^{n-2})\sharp(Z_{l'-1} \bmod 2^{r+l'}) + 3 \cdot 2^{n-2} \sum_{i=3}^{l'-2} \sharp(Z_i \bmod 2^{r+i+1}) \\ &+ 7 \cdot 2^{n-3}\sharp(Z_1 \bmod 2^{r+3}) + 2^{n-3}\sharp(Z_0 \bmod 2^{r+3}).\end{aligned}$$

By Corollary 5.17 and Lemma 5.3, we have

$$\sharp(Z_i \bmod 2^{r+i+1}) \leq 2^i$$

for $3 \leq i \leq l' - 1$. We have seen

$$\sharp(Z_1 \bmod 2^{r+3}) \leq 2^2$$

before. Consequently, we get

$$\begin{aligned}\sharp Z_0 &\leq (2^{2(n-l')} - 2^{n-2}) \cdot 2^{l'-1} + 3 \cdot 2^{n-2} \sum_{i=3}^{l'-2} 2^i + 7 \cdot 2^{n-3} \cdot 2^2 + 2^{n-3}\sharp(Z_0 \bmod 2^{r+3}) \\ &= b(u, 2)_n + 2^{n-3}(\sharp(H/H_{r+3}) \cap \mathrm{Conj}(u^{2^r}) - 4).\end{aligned}$$

□

7. Proof of the main theorem

Now we prove Theorem 3.9. Recall the four types of groups

$$\begin{aligned}B &= \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \right\} \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}), \\ C &= \left\{ \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}, \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix} \right\} \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}), \\ D &= \left\{ \begin{pmatrix} x & y \\ \lambda y & x \end{pmatrix}, \begin{pmatrix} x & y \\ -\lambda y & -x \end{pmatrix} \right\} \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}), \\ E &= (\text{an exceptional subgroup}) \cap \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}).\end{aligned}$$

For a subgroup $H \subseteq G = \mathrm{SL}_2(\mathbb{Z}/p^n\mathbb{Z})$, recall (4.1):

$$\delta_H = 1 - 3\frac{\sharp H \cap \mathrm{Conj}(\sigma)}{\sharp \mathrm{Conj}(\sigma)} - 4\frac{\sharp H \cap \mathrm{Conj}(\tau)}{\sharp \mathrm{Conj}(\tau)} - 6\frac{\sharp I\backslash G/H}{[G : H]},$$

and (4.2), (4.3):

$$\frac{\sharp I\backslash G/H}{[G : H]} = \sum_{s=0}^{n-1} \frac{p-1}{p^{s+1}} \frac{\sharp H \cap \mathrm{Conj}(u^{p^s})}{\sharp \mathrm{Conj}(u^{p^s})} + \frac{1}{p^n} \leq \sum_{s=0}^{t-1} \frac{p-1}{p^{s+1}} \frac{\sharp H \cap \mathrm{Conj}(u^{p^s})}{\sharp \mathrm{Conj}(u^{p^s})} + \frac{1}{p^t},$$

where $1 \leq t \leq n$.

Lemma 7.1. *Let $p \geq 5$ and assume a subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ is contained in an exceptional subgroup E. If $p \geq 17$, then $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$, we have $\sharp\mathrm{Conj}(\sigma) \geq (p-1)p$ and $\sharp\mathrm{Conj}(\tau) \geq (p-1)p$ by Lemma 5.1. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z})$ we have $\sharp E \cap \mathrm{Conj}(\sigma) \leq 30$, $\sharp E \cap \mathrm{Conj}(\tau) \leq 20$ and $\sharp E \cap \mathrm{Conj}(u) = 0$. Since $\sharp E \cap \mathrm{Conj}(u) = 0$, we have $\frac{\sharp I\backslash G/H}{[G:H]} = \frac{1}{p}$. Therefore $\delta_H \geq 1 - 3 \cdot \frac{30}{(p-1)p} - 4 \cdot \frac{20}{(p-1)p} - 6 \cdot \frac{1}{p} = \frac{p^2-7p-164}{(p-1)p} > 0$ if $p \geq 17$.

□

Proposition 7.2. *Assume $p = 19$. For any slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/19^2\mathbb{Z})$ satisfying $H/H_1 \subseteq B$, we have $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/19^2\mathbb{Z})$, we have $\sharp\mathrm{Conj}(\tau) = 20 \cdot 19^3$ and $\sharp\mathrm{Conj}(u) = \frac{1}{2}(p^2-1)p^2$ by Lemma 5.1. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/19\mathbb{Z})$ we have $\sharp B \cap \mathrm{Conj}(\sigma) = 0$, $\sharp B \cap \mathrm{Conj}(\tau) = 2p = 38$ and $\sharp B \cap \mathrm{Conj}(u) = \frac{1}{2}(p-1) = 9$. By Corollary 6.10, we have $\sharp H \cap \mathrm{Conj}(\tau) \leq a(\tau, p)_2 + p(38-2) = 74 \cdot 19$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} \leq \frac{74\cdot 19}{20\cdot 19^3} = \frac{37}{10\cdot 19^2}$. We have $\sharp H \cap \mathrm{Conj}(u) \leq p^2\sharp(H/H_1)\cap\mathrm{Conj}(u)$ by Lemma 5.3. Thus $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)} \leq \frac{p^2\cdot\frac{1}{2}(p-1)}{\frac{1}{2}(p^2-1)p^2} = \frac{1}{p+1}$. Therefore $\frac{\sharp I\backslash G/H}{[G:H]} \leq \frac{p-1}{p}\cdot\frac{1}{p+1} + \frac{1}{p} = \frac{2}{p+1} = \frac{1}{10}$. Consequently, $\delta_H \geq 1 - 3\cdot 0 - 4\cdot\frac{37}{10\cdot 19^2} - 6\cdot\frac{1}{10} = \frac{1805-74-1083}{5\cdot 19^2} > 0$, as required.

□

Proposition 7.3. *Assume $p = 17$. For any slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/17^2\mathbb{Z})$ satisfying $H/H_1 \subseteq B$, we have $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/17^2\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma) = 18\cdot 17^3$ and $\sharp\mathrm{Conj}(u) = \frac{1}{2}(p^2-1)p^2$ by Lemma 5.1. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/17\mathbb{Z})$ we have $\sharp B\cap\mathrm{Conj}(\sigma) = 2p = 34$, $\sharp B \cap \mathrm{Conj}(\tau) = 0$ and $\sharp B \cap \mathrm{Conj}(u) = \frac{1}{2}(p-1) = 8$. By Corollary 6.9, we have $\sharp H \cap \mathrm{Conj}(\sigma) \leq a(\sigma, p)_2 + p(34-2) = 66\cdot 17$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)} \leq \frac{66\cdot 17}{18\cdot 17^3} = \frac{11}{3\cdot 17^2}$. We also have $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)} \leq \frac{1}{p+1}$ by the same reason as in Proposition 7.2. Thus $\frac{I\backslash G/H}{[G:H]} \leq \frac{p-1}{p}\cdot\frac{1}{p+1} + \frac{1}{p} = \frac{2}{p+1} = \frac{1}{9}$. Consequently, $\delta_H \geq 1 - 3\cdot\frac{11}{3\cdot 17^2} - 4\cdot 0 - 6\cdot\frac{1}{9} = \frac{867-33-578}{3\cdot 17^2} > 0$.

□

Proposition 7.4. *Assume $p = 13$. Take a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/13^2\mathbb{Z})$. If H/H_1 is contained in B or E, then $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/13^2\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma) = \sharp\mathrm{Conj}(\tau) = 14\cdot 13^3$, $\sharp\mathrm{Conj}(u) = \frac{1}{2}(p^2-1)p^2$ and $\sharp\mathrm{Conj}(u^p) = \frac{1}{2}(p^2-1)$ by Lemma 5.1, 5.2.

Suppose $H/H_1 \subseteq B$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/13\mathbb{Z})$ we have $\sharp B \cap \mathrm{Conj}(\sigma) = 2p = 26$, $\sharp B \cap \mathrm{Conj}(\tau) = 2p = 26$ and $\sharp B \cap \mathrm{Conj}(u) = \frac{1}{2}(p-1) = 6$. By Corollary 6.9, we have $\sharp H \cap \mathrm{Conj}(\sigma) \leq a(\sigma, p)_2 + p(26-2) = 50\cdot 13$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)} \leq \frac{50\cdot 13}{14\cdot 13^3} = \frac{25}{7\cdot 13^2}$. The same calculation shows $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} \leq \frac{25}{7\cdot 13^2}$. We have $B \cap \mathrm{Conj}(u) = \left\{\begin{pmatrix} 1 & s^2 \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}/p\mathbb{Z}) | (p, s) = 1\right\}$ by Lemma 4.9, and $V_{u'} = V_u = \left\{1 + p\begin{pmatrix} a & b \\ 0 & -a \end{pmatrix}\right\}$ for any $u' \in \left\{\begin{pmatrix} 1 & s^2 \\ 0 & 1 \end{pmatrix} | (p, s) = 1\right\}$ by Lemma 5.3 and Corollary 5.16.

If H contains V_u, then the $(2,1)$ entry of an element of $H \cap \mathrm{Conj}(u^p)$ must be zero since H is slim, thus $H \cap \mathrm{Conj}(u^p) \subseteq \left\{\begin{pmatrix} 1 & ps^2 \\ 0 & 1 \end{pmatrix} | (p, s) = 1\right\}$. Hence $\frac{\sharp H\cap\mathrm{Conj}(u^p)}{\sharp\mathrm{Conj}(u^p)} \leq \frac{\frac{1}{2}(p-1)}{\frac{1}{2}(p^2-1)} = \frac{1}{p+1}$. We also have $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)} \leq \frac{1}{p+1}$ as in Proposition 7.2. Thus $\frac{\sharp I\backslash G/H}{[G:H]} \leq \frac{p-1}{p}\cdot\frac{1}{p+1} + \frac{p-1}{p^2}\cdot\frac{1}{p+1} + \frac{1}{p^2} = \frac{1}{p} = \frac{1}{13}$. Therefore $\delta_H \geq 1 - 3\cdot\frac{25}{7\cdot 13^2} - 4\cdot\frac{25}{7\cdot 13^2} - 6\cdot\frac{1}{13} = \frac{1183-75-100-546}{7\cdot 13^2} > 0$.

If H does not contain V_u, for each element $x \in (H/H_1)\cap\mathrm{Conj}(u) \subseteq B\cap\mathrm{Conj}(u)$ (there are at most $\frac{1}{2}(p-1)$ such elements by Lemma 4.9) we have $\sharp(f_{2,1}^{H,u})^{-1}(x) \leq p = 13$ by Lemma 6.5. Hence $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)} \leq \frac{p\cdot\frac{1}{2}(p-1)}{\frac{1}{2}(p^2-1)p^2} = \frac{1}{p(p+1)}$ and $\frac{\sharp I\backslash G/H}{[G:H]} \leq$

$\frac{p-1}{p}\cdot\frac{1}{p(p+1)}+\frac{1}{p}=\frac{p^2+2p-1}{(p+1)p^2}=\frac{97}{7\cdot 13^2}$. Therefore $\delta_H \geq 1-3\cdot\frac{25}{7\cdot 13^2}-4\cdot\frac{25}{7\cdot 13^2}-6\cdot\frac{97}{7\cdot 13^2}=\frac{1183-75-100-582}{7\cdot 13^2}>0$.

Next suppose $H/H_1 \subseteq E$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/13\mathbb{Z})$ we have $\sharp E\cap \mathrm{Conj}(\sigma)\leq 18$, $\sharp E\cap\mathrm{Conj}(\tau)\leq 8$ and $\sharp E\cap\mathrm{Conj}(u)=0$. By Corollary 6.9, we have $\sharp H\cap\mathrm{Conj}(\sigma)\leq a(\sigma,p)_2+p(18-2)=42\cdot 13$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)}\leq\frac{42\cdot 13}{14\cdot 13^3}=\frac{3}{13^2}$. By Corollary 6.10, we have $\sharp H\cap\mathrm{Conj}(\tau)\leq a(\tau,p)_2+p(8-2)=32\cdot 13$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)}\leq\frac{32\cdot 13}{14\cdot 13^3}=\frac{16}{7\cdot 13^2}$. Since $\sharp(H/H_1)\cap\mathrm{Conj}(u)=0$, we have $\frac{\sharp I\backslash G/H}{[G:H]}\leq\frac{1}{p}=\frac{1}{13}$ (take $t=1$ in (4.3)). Therefore $\delta_H\geq 1-3\cdot\frac{3}{13^2}-4\cdot\frac{16}{7\cdot 13^2}-6\cdot\frac{1}{13}=\frac{1183-63-64-546}{7\cdot 13^2}>0$.

□

Proposition 7.5. *Assume $p=11$. Take a slim subgroup $H\subseteq\mathrm{SL}_2(\mathbb{Z}/11^2\mathbb{Z})$. If H/H_1 is contained in B, D or E, then $\delta_H>0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/11^2\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma)=\sharp\mathrm{Conj}(\tau)=10\cdot 11^3$, $\sharp\mathrm{Conj}(u)=\frac{1}{2}(p^2-1)p^2$ and $\sharp\mathrm{Conj}(u^p)=\frac{1}{2}(p^2-1)$ by Lemma 5.1, 5.2.

Suppose $H/H_1\subseteq B$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/11\mathbb{Z})$ we have $\sharp B\cap\mathrm{Conj}(\sigma)=0$, $\sharp B\cap\mathrm{Conj}(\tau)=0$ and $\sharp B\cap\mathrm{Conj}(u)=\frac{1}{2}(p-1)=5$.

If H contains V_u, then the same calculation as in Proposition 7.4 shows $\frac{\sharp I\backslash G/H}{[G:H]}\leq\frac{1}{p}=\frac{1}{11}$. Therefore $\delta_H\geq 1-3\cdot 0-4\cdot 0-6\cdot\frac{1}{11}=\frac{11-6}{11}>0$.

If H does not contain V_u, as in Proposition 7.4, $\frac{\sharp I\backslash G/H}{[G:H]}\leq\frac{p^2+2p-1}{(p+1)p^2}=\frac{71}{2\cdot 3\cdot 11^2}$. Therefore $\delta_H\geq 1-3\cdot 0-4\cdot 0-6\cdot\frac{71}{2\cdot 3\cdot 11^2}=\frac{121-71}{11^2}>0$.

Next suppose $H/H_1\subseteq E$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/11\mathbb{Z})$ we have $\sharp E\cap\mathrm{Conj}(\sigma)\leq 30$, $\sharp E\cap\mathrm{Conj}(\tau)\leq 20$ and $\sharp E\cap\mathrm{Conj}(u)=0$. By Corollary 6.9, we have $\sharp H\cap\mathrm{Conj}(\sigma)\leq a(\sigma,p)_2+p(30-2)=50\cdot 11$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)}\leq\frac{50\cdot 11}{10\cdot 11^3}=\frac{5}{11^2}$. By Corollary 6.10, we have $\sharp H\cap\mathrm{Conj}(\tau)\leq a(\tau,p)_2+p(20-2)=40\cdot 11$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)}\leq\frac{40\cdot 11}{10\cdot 11^3}=\frac{4}{11^2}$. Since $\sharp(H/H_1)\cap\mathrm{Conj}(u)=0$, we have $\frac{\sharp I\backslash G/H}{[G:H]}\leq\frac{1}{p}=\frac{1}{11}$. Therefore $\delta_H\geq 1-3\cdot\frac{5}{11^2}-4\cdot\frac{4}{11^2}-6\cdot\frac{1}{11}=\frac{121-15-16-66}{11^2}>0$.

Finally, suppose $H/H_1\subseteq D$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/11\mathbb{Z})$ we have $\sharp D\cap\mathrm{Conj}(\sigma)=p+3=14$, $\sharp D\cap\mathrm{Conj}(\tau)=2$ and $\sharp D\cap\mathrm{Conj}(u)=0$. Since $\sharp D\cap\mathrm{Conj}(\sigma)<30$, $\sharp D\cap\mathrm{Conj}(\tau)<20$ and $\sharp D\cap\mathrm{Conj}(u)=0$, the calculation for E shows $\delta_H>0$ in this case.

□

Proposition 7.6. *Assume $p=7$. Take a slim subgroup $H\subseteq\mathrm{SL}_2(\mathbb{Z}/7^3\mathbb{Z})$. If H/H_1 is contained in B, C, D or E , then $\delta_H>0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/7^3\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma)=6\cdot 7^5$, $\sharp\mathrm{Conj}(\tau)=8\cdot 7^5$, $\sharp\mathrm{Conj}(u)=\frac{1}{2}(p^2-1)p^4$, $\sharp\mathrm{Conj}(u^p)=\frac{1}{2}(p^2-1)p^2$ and $\sharp\mathrm{Conj}(u^{p^2})=\frac{1}{2}(p^2-1)$ by Lemma 5.1, 5.2.

Suppose $H/H_1\subseteq B$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/7\mathbb{Z})$ we have $\sharp B\cap\mathrm{Conj}(\sigma)=0$, $\sharp B\cap\mathrm{Conj}(\tau)=2p=14$ and $\sharp B\cap\mathrm{Conj}(u)=\frac{1}{2}(p-1)=3$. By Corollary 6.10, we have $\sharp H\cap\mathrm{Conj}(\tau)\leq a(\tau,p)_3+p^2(14-2)=110\cdot 7^2$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)}\leq\frac{110\cdot 7^2}{8\cdot 7^5}=\frac{55}{2^2\cdot 7^3}$.

If H/H_2 contains $V_u^{2,1}=\left\{1+p\begin{pmatrix}a & b\\ 0 & -a\end{pmatrix}\right\}$, then the $(2,1)$ entry of an element of $(H/H_2)\cap\mathrm{Conj}(u^p)$ must be zero as we have seen in Proposition 7.4, thus $(H/H_2) \cap \mathrm{Conj}(u^p) \subseteq \left\{ \begin{pmatrix} 1 & ps^2 \\ 0 & 1 \end{pmatrix} | (p,s) = 1 \right\}$. We have $\sharp H \cap \mathrm{Conj}(u^p) \leq p^2 \sharp (H/H_2) \cap \mathrm{Conj}(u^p)$ by Lemma 5.3. Hence $\frac{\sharp H \cap \mathrm{Conj}(u^p)}{\sharp \mathrm{Conj}(u^p)} \leq \frac{p^2 \cdot \frac{1}{2}(p-1)}{\frac{1}{2}(p^2-1)p^2} = \frac{1}{p+1}$. As H/H_2 contains $\left\{ 1 + p \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} \right\}$, the subgroup H contains $\left\{ 1 + p^2 \begin{pmatrix} a & b \\ 0 & -a \end{pmatrix} \right\}$. Thus the $(2,1)$ entry of an element of $H \cap \mathrm{Conj}(u^{p^2})$ must be zero since H is slim, and $H \cap \mathrm{Conj}(u^{p^2}) \subseteq \left\{ \begin{pmatrix} 1 & p^2 s^2 \\ 0 & 1 \end{pmatrix} | (p,s) = 1 \right\}$. Hence $\frac{\sharp H \cap \mathrm{Conj}(u^{p^2})}{\sharp \mathrm{Conj}(u^{p^2})} \leq \frac{\frac{1}{2}(p-1)}{\frac{1}{2}(p^2-1)} = \frac{1}{p+1}$. We also have $\frac{\sharp H \cap \mathrm{Conj}(u)}{\sharp \mathrm{Conj}(u)} \leq \frac{1}{p+1}$ as in Proposition 7.2. Therefore $\frac{\sharp I \backslash G/H}{[G:H]} \leq \frac{p-1}{p} \cdot \frac{1}{p+1} + \frac{p-1}{p^2} \cdot \frac{1}{p+1} + \frac{p-1}{p^3} \cdot \frac{1}{p+1} + \frac{1}{p^3} = \frac{p^2+1}{(p+1)p^2} = \frac{25}{2^2 \cdot 7^2}$. Consequently, $\delta_H \geq 1 - 3 \cdot 0 - 4 \cdot \frac{55}{2^2 \cdot 7^3} - 6 \cdot \frac{25}{2^2 \cdot 7^2} = \frac{686-110-525}{2 \cdot 7^3} > 0$.

If H/H_2 does not contain $V_u^{2,1}$, for each element $x \in (H/H_1) \cap \mathrm{Conj}(u) \subseteq B \cap \mathrm{Conj}(u)$ (there are at most $\frac{1}{2}(p-1)$ such elements by Lemma 4.9) we have $\sharp (f_{3,1}^{H,u})^{-1}(x) \leq p^2 \sharp (f_{2,1}^{H,u})^{-1}(x) \leq p^3$ by Lemma 5.3, 6.5. Thus $\frac{\sharp H \cap \mathrm{Conj}(u)}{\sharp \mathrm{Conj}(u)} \leq \frac{p^3 \cdot \frac{1}{2}(p-1)}{\frac{1}{2}(p^2-1)p^4} = \frac{1}{(p+1)p} = \frac{1}{2^3 \cdot 7}$. By Corollary 6.12, we have $\sharp H \cap \mathrm{Conj}(u^p) \leq a(u,p)_2 + p(\frac{1}{2}(p^2-1) - \frac{1}{2}(p-1)) = (p-1)p^2$. Thus $\frac{\sharp H \cap \mathrm{Conj}(u^p)}{\sharp \mathrm{Conj}(u^p)} \leq \frac{(p-1)p^2}{\frac{1}{2}(p^2-1)p^2} = \frac{2}{p+1} = \frac{1}{4}$. Therefore $\frac{\sharp I \backslash G/H}{[G:H]} \leq \frac{p-1}{p} \cdot \frac{1}{(p+1)p} + \frac{p-1}{p^2} \cdot \frac{2}{p+1} + \frac{1}{p^2} = \frac{4p-2}{(p+1)p^2} = \frac{13}{2^2 \cdot 7^2}$. Consequently, $\delta_H \geq 1 - 3 \cdot 0 - 4 \cdot \frac{55}{2^2 \cdot 7^3} - 6 \cdot \frac{13}{2^2 \cdot 7^2} = \frac{686-110-273}{2 \cdot 7^3} > 0$.

Next suppose $H/H_1 \subseteq E$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/7\mathbb{Z})$ we have $\sharp E \cap \mathrm{Conj}(\sigma) \leq 18$, $\sharp E \cap \mathrm{Conj}(\tau) \leq 8$ and $\sharp E \cap \mathrm{Conj}(u) = 0$. By Corollary 6.9, we have $\sharp H \cap \mathrm{Conj}(\sigma) \leq a(\sigma,p)_3 + p^2(18-2) = 114 \cdot 7^2$. Hence $\frac{\sharp H \cap \mathrm{Conj}(\sigma)}{\sharp \mathrm{Conj}(\sigma)} \leq \frac{114 \cdot 7^2}{6 \cdot 7^5} = \frac{19}{7^3}$. By Corollary 6.10, we have $\sharp H \cap \mathrm{Conj}(\tau) \leq a(\tau,p)_3 + p^2(8-2) = 104 \cdot 7^2$. Hence $\frac{\sharp H \cap \mathrm{Conj}(\tau)}{\sharp \mathrm{Conj}(\tau)} \leq \frac{104 \cdot 7^2}{8 \cdot 7^5} = \frac{13}{7^3}$. The same calculation as above (the case that $H/H_1 \subseteq B$ and H/H_2 does not contain $V_u^{2,1}$) shows the inequality $\frac{\sharp H \cap \mathrm{Conj}(u^p)}{\sharp \mathrm{Conj}(u^p)} \leq \frac{2}{p+1} = \frac{1}{4}$. Therefore $\frac{\sharp I \backslash G/H}{[G:H]} \leq \frac{p-1}{p} \cdot 0 + \frac{p-1}{p^2} \cdot \frac{2}{p+1} + \frac{1}{p^2} = \frac{3p-1}{(p+1)p^2} = \frac{5}{2 \cdot 7^2}$. Consequently, $\delta_H \geq 1 - 3 \cdot \frac{19}{7^3} - 4 \cdot \frac{13}{7^3} - 6 \cdot \frac{5}{2 \cdot 7^2} = \frac{343-57-52-105}{7^3} > 0$.

Suppose $H/H_1 \subseteq C$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/7\mathbb{Z})$ we have $\sharp C \cap \mathrm{Conj}(\sigma) = p-1 = 6 < 18$, $\sharp C \cap \mathrm{Conj}(\tau) = 2 < 8$, $\sharp C \cap \mathrm{Conj}(u) = 0$. The calculation for E shows $\delta_H > 0$ in this case.

Finally, suppose $H/H_1 \subseteq D$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/7\mathbb{Z})$ we have $\sharp D \cap \mathrm{Conj}(\sigma) = p+3 = 10 < 18$, $\sharp D \cap \mathrm{Conj}(\tau) = 0 < 8$ and $\sharp D \cap \mathrm{Conj}(u) = 0$. The calculation for E also shows $\delta_H > 0$ in this case.

□

Proposition 7.7. *Assume $p = 5$. Take a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/5^3\mathbb{Z})$. If H/H_1 is contained in C, then $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/5^3\mathbb{Z})$ we have $\sharp \mathrm{Conj}(\sigma) = 6 \cdot 5^5$, and $\sharp \mathrm{Conj}(u^p) = \frac{1}{2}(5^2-1)5^2 = 12 \cdot 5^2$ by Lemma 5.1, 5.2. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/5\mathbb{Z})$ we have $\sharp C \cap \mathrm{Conj}(\sigma) = p+1 = 6$, $\sharp C \cap \mathrm{Conj}(\tau) = 0$ and $\sharp C \cap \mathrm{Conj}(u) = 0$. By Corollary 6.9, we have $\sharp H \cap \mathrm{Conj}(\sigma) \leq a(\sigma,p)_3 + p^3(6-2) = 54 \cdot 5^2$. Hence $\frac{\sharp H \cap \mathrm{Conj}(\sigma)}{\sharp \mathrm{Conj}(\sigma)} \leq \frac{54 \cdot 5^2}{6 \cdot 5^5} = \frac{9}{5^3}$. In $\mathrm{SL}_2(\mathbb{Z}/5^2\mathbb{Z})$ we have $\sharp \mathrm{Conj}(u^5) = 12$ by Lemma 5.2. By Corollary 6.12, we have $\sharp H \cap \mathrm{Conj}(u^p) \leq a(u,p)_2 + p(12 - \frac{1}{2}(5-1)) = 4 \cdot 5^2$. Hence $\frac{\sharp H \cap \mathrm{Conj}(u^p)}{\sharp \mathrm{Conj}(u^p)} \leq$

$\frac{4\cdot 5^2}{12\cdot 5^2} = \frac{1}{3}$. Therefore $\frac{\sharp I\backslash G/H}{[G:H]} \leq \frac{5-1}{5}\cdot 0 + \frac{5-1}{5^2}\cdot \frac{1}{3} + \frac{1}{5^2} = \frac{7}{3\cdot 5^2}$. Consequently, $\delta_H \geq 1 - 3\cdot \frac{9}{5^3} - 4\cdot 0 - 6\cdot \frac{7}{3\cdot 5^2} = \frac{125-27-70}{5^3} > 0$.

□

Proposition 7.8. *Assume $p = 5$. Take a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/5^4\mathbb{Z})$. If H/H_1 is contained in B, D or E, then $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/5^4\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma) = 6\cdot 5^7$, $\sharp\mathrm{Conj}(\tau) = 4\cdot 5^7$, $\sharp\mathrm{Conj}(u) = 12\cdot 5^6$, $\sharp\mathrm{Conj}(u^p) = 12\cdot 5^4$ and $\sharp\mathrm{Conj}(u^{p^2}) = 12\cdot 5^2$ by Lemma 5.1, 5.2.

Suppose $H/H_1 \subseteq B$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/5\mathbb{Z})$ we have $\sharp B\cap\mathrm{Conj}(\sigma) = 2p = 10$, $\sharp B\cap\mathrm{Conj}(\tau) = 0$ and $\sharp B\cap\mathrm{Conj}(u) = \frac{1}{2}(p-1) = 2$. By Corollary 6.9, we have $\sharp H\cap\mathrm{Conj}(\sigma) \leq a(\sigma,p)_4 + p^3(10-2) = 66\cdot 5^3$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)} \leq \frac{66\cdot 5^3}{6\cdot 5^7} = \frac{11}{5^4}$. By Corollary 6.12, we have $\sharp H\cap\mathrm{Conj}(u) \leq a(u,p)_4 = 18\cdot 5^4$. Hence $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)} \leq \frac{18\cdot 5^4}{12\cdot 5^6} = \frac{3}{2\cdot 5^2}$. In $\mathrm{SL}_2(\mathbb{Z}/5^2\mathbb{Z})$ we have $\sharp\mathrm{Conj}(u^p) = 12$ by Lemma 5.2. Again by Corollary 6.12, we have $\sharp H\cap\mathrm{Conj}(u^p) \leq a(u,p)_3 + p^2(12-\frac{1}{2}(5-1)) = 12\cdot 5^3$. Hence $\frac{\sharp H\cap\mathrm{Conj}(u^p)}{\sharp\mathrm{Conj}(u^p)} \leq \frac{12\cdot 5^3}{12\cdot 5^4} = \frac{1}{5}$. Similarly $\sharp H\cap\mathrm{Conj}(u^{p^2}) \leq a(u,p)_2 + p(12-\frac{1}{2}(5-1)) = 4\cdot 5^2$. Hence $\frac{\sharp H\cap\mathrm{Conj}(u^{p^2})}{\sharp\mathrm{Conj}(u^{p^2})} \leq \frac{4\cdot 5^2}{12\cdot 5^2} = \frac{1}{3}$. Therefore $\frac{\sharp I\backslash G/H}{[G:H]} \leq \frac{5-1}{5}\cdot\frac{3}{2\cdot 5^2} + \frac{5-1}{5^2}\cdot\frac{1}{5} + \frac{5-1}{5^3}\cdot\frac{1}{3} + \frac{1}{5^3} = \frac{37}{3\cdot 5^3}$. Consequently, $\delta_H \geq 1 - 3\cdot\frac{11}{5^4} - 4\cdot 0 - 6\cdot\frac{37}{3\cdot 5^3} = \frac{625-33-370}{5^4} > 0$.

Next suppose $H/H_1 \subseteq E$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/5\mathbb{Z})$ we have $\sharp E\cap\mathrm{Conj}(\sigma) \leq 18$, $\sharp E\cap\mathrm{Conj}(\tau) \leq 8$ and $\sharp E\cap\mathrm{Conj}(u) = 0$. By Corollary 6.9, we have $\sharp H\cap\mathrm{Conj}(\sigma) \leq a(\sigma,p)_4 + p^3(18-2) = 74\cdot 5^3$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)} \leq \frac{74\cdot 5^3}{6\cdot 5^7} = \frac{37}{3\cdot 5^4}$. By Corollary 6.10, we have $\sharp H\cap\mathrm{Conj}(\tau) \leq a(\tau,p)_4 + p^3(8-2) = 64\cdot 5^3$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} \leq \frac{64\cdot 5^3}{4\cdot 5^7} = \frac{16}{5^4}$. The calculation in the Borel case shows $\frac{\sharp I\backslash G/H}{[G:H]} \leq \frac{5-1}{5}\cdot 0 + \frac{5-1}{5^2}\cdot\frac{1}{5} + \frac{5-1}{5^3}\cdot\frac{1}{3} + \frac{1}{5^3} = \frac{19}{3\cdot 5^3}$. Consequently, $\delta_H \geq 1 - 3\cdot\frac{37}{3\cdot 5^4} - 4\cdot\frac{16}{5^4} - 6\cdot\frac{19}{3\cdot 5^3} = \frac{625-37-64-190}{5^4} > 0$.

Finally, suppose $H/H_1 \subseteq D$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/5\mathbb{Z})$ we have $\sharp D\cap\mathrm{Conj}(\sigma) = p+1 = 6 < 18$, $\sharp D\cap\mathrm{Conj}(\tau) = 2 < 8$ and $\sharp D\cap\mathrm{Conj}(u) = 0$. The calculation for E shows $\delta_H > 0$ in this case.

□

Proposition 7.9. *Assume $p = 3$. Take a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/3^6\mathbb{Z})$. If H/H_1 is contained in B, C, D or $H/H_1 = \mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$, then $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/3^6\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma) = 2\cdot 3^{11}$, $\sharp\mathrm{Conj}(\tau) = 4\cdot 3^{10}$ and $\sharp\mathrm{Conj}(u^{3^i}) = 4\cdot 3^{10-2i}$ for $0 \leq i \leq 4$ by Lemma 5.1, 5.2.

Suppose $H/H_1 \subseteq B$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$ we have $\sharp B\cap\mathrm{Conj}(\sigma) = 0$, $\sharp B\cap\mathrm{Conj}(\tau) = 1$ and $\sharp B\cap\mathrm{Conj}(u) = 1$. By Corollary 6.11, we have $\sharp H\cap\mathrm{Conj}(\tau) \leq a(\tau,3)_6 = 13\cdot 3^6$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} \leq \frac{13\cdot 3^6}{4\cdot 3^{10}} = \frac{13}{2^2\cdot 3^4}$. By Corollary 6.12, we have $\sharp H\cap\mathrm{Conj}(u) \leq a(u,3)_6 = 17\cdot 3^6$. Hence $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)} \leq \frac{17\cdot 3^6}{4\cdot 3^{10}} = \frac{17}{2^2\cdot 3^4}$. In $\mathrm{SL}_2(\mathbb{Z}/3^{r+1}\mathbb{Z})$ we have $\sharp\mathrm{Conj}(u^{3^r}) = \frac{1}{2}(3^2-1) = 4$ by Lemma 5.2. Again by Corollary 6.12, we have $\sharp H\cap\mathrm{Conj}(u^{3^i}) \leq a(u,3)_{6-i} + 3^{5-i}(4-\frac{1}{2}(3-1)) = a(u,3)_{6-i} + 3^{6-i}$ for $1 \leq i \leq 4$. Hence

$$\frac{\sharp H\cap\mathrm{Conj}(u^{3^i})}{\sharp\mathrm{Conj}(u^{3^i})} \leq \frac{a(u,3)_{6-i} + 3^{6-i}}{4\cdot 3^{10-2i}}$$

for $1 \leq i \leq 4$. Therefore

$$\begin{aligned}\frac{\sharp I\backslash G/H}{[G:H]} \leq & \frac{3-1}{3}\cdot\frac{17}{2^2\cdot 3^4}+\sum_{i=1}^{4}\frac{3-1}{3^{i+1}}\cdot\frac{a(u,3)_{6-i}+3^{6-i}}{4\cdot 3^{10-2i}}+\frac{1}{3^5}\\ = & \frac{17+12+6+4+2+2}{2\cdot 3^5}=\frac{43}{2\cdot 3^5}.\end{aligned}$$

Consequently, $\delta_H \geq 1-3\cdot 0-4\cdot\frac{13}{2^2\cdot 3^4}-6\cdot\frac{43}{2\cdot 3^5}=\frac{81-13-43}{3^4}>0$.

Next suppose $H/H_1=\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$. Since $H/H_2 \subsetneq \mathrm{SL}_2(\mathbb{Z}/3^2\mathbb{Z})$ by Lemma 2.1, we have $H\cap\mathrm{Conj}(u)=\emptyset$ by Lemma 4.4. Lemma 5.1 shows that in $\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma)=6$ and $\sharp\mathrm{Conj}(\tau)=4$. By Corollary 6.9, we have $\sharp H\cap\mathrm{Conj}(\sigma)\leq a(\sigma,3)_6+3^5(6-2)=14\cdot 3^6$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)}\leq\frac{14\cdot 3^6}{2\cdot 3^{11}}=\frac{7}{3^5}$. By Corollary 6.11, we have $\sharp H\cap\mathrm{Conj}(\tau)\leq a(\tau,3)_6+3^5(4-1)=14\cdot 3^6$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)}\leq\frac{14\cdot 3^6}{4\cdot 3^{10}}=\frac{7}{2\cdot 3^4}$. The same calculation as in the Borel case shows

$$\frac{\sharp I\backslash G/H}{[G:H]}\leq\frac{3-1}{3}\cdot 0+\sum_{i=1}^{4}\frac{3-1}{3^{i+1}}\cdot\frac{a(u,3)_{6-i}+3^{6-i}}{4\cdot 3^{10-2i}}+\frac{1}{3^5}=\frac{13}{3^5}.$$

Consequently, $\delta_H\geq 1-3\cdot\frac{7}{3^5}-4\cdot\frac{7}{2\cdot 3^4}-6\cdot\frac{13}{3^5}=\frac{81-7-14-26}{3^4}>0$.

Suppose $H/H_1\subseteq C$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$ we have $\sharp C\cap\mathrm{Conj}(\sigma)=2<6$, $\sharp C\cap\mathrm{Conj}(\tau)=0<4$, $\sharp C\cap\mathrm{Conj}(u)=0$. The calculation for the case $H/H_1=\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$ shows $\delta_H>0$ in this case.

Finally, suppose $H/H_1\subseteq D$. Lemma 4.9 shows that in $\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$ we have $\sharp D\cap\mathrm{Conj}(\sigma)=6$, $\sharp D\cap\mathrm{Conj}(\tau)=0<4$ and $\sharp D\cap\mathrm{Conj}(u)=0$. The calculation for the case $H/H_1=\mathrm{SL}_2(\mathbb{Z}/3\mathbb{Z})$ also shows $\delta_H>0$ in this case.

□

As is well-known, we have $\mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})\cong S_3$ as a group. All non-trivial subgroups of $\mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$ are the following:

$$\begin{aligned}B=&\left\{\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}1&1\\0&1\end{pmatrix}\right\},\left\{\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}1&0\\1&1\end{pmatrix}\right\},\left\{\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}0&1\\1&0\end{pmatrix}\right\},\\ F=&\left\{\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}1&1\\1&0\end{pmatrix},\begin{pmatrix}0&1\\1&1\end{pmatrix}\right\}.\end{aligned}$$

The first three groups are conjugate.

Proposition 7.10. *Assume $p=2$. Take a slim subgroup $H\subseteq\mathrm{SL}_2(\mathbb{Z}/2^{11}\mathbb{Z})$. If H/H_1 is contained in B, then $\delta_H>0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/2^{11}\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma)=3\cdot 2^{19}$ and $\sharp\mathrm{Conj}(u^{2^i})=3\cdot 2^{18-2i}$ for $0\leq i\leq 7$ by Lemma 5.1, 5.2. By Lemma 4.6, we see that in $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ we have $\sharp f_{2,1}^{-1}(B)\cap\mathrm{Conj}(\sigma)=2$, $\sharp f_{2,1}^{-1}(B)\cap\mathrm{Conj}(\tau)=0$, $\sharp f_{2,1}^{-1}(B)\cap\mathrm{Conj}(u)=2$ and $\sharp f_{2,1}^{-1}(B)\cap\mathrm{Conj}(u^2)=3$. By Proposition 6.14, we have $\sharp H\cap\mathrm{Conj}(\sigma)\leq a(\sigma,2)_{11}=11\cdot 2^{12}$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)}\leq\frac{11\cdot 2^{12}}{3\cdot 2^{19}}=\frac{11}{3\cdot 2^7}$. We have $\sharp(H/H_3)\cap\mathrm{Conj}(u)\leq\sharp f_{3,1}^{-1}(B)\cap\mathrm{Conj}(u)=4$ by Remark 5.4. By Corollary 6.13, we have $\sharp H\cap\mathrm{Conj}(u)\leq a(u,2)_{11}+2^{10}(4-2)=23\cdot 2^{11}$. Hence $\frac{\sharp H\cap\mathrm{Conj}(u)}{\sharp\mathrm{Conj}(u)}\leq\frac{23\cdot 2^{11}}{3\cdot 2^{18}}=\frac{23}{3\cdot 2^7}$. In $\mathrm{SL}_2(\mathbb{Z}/2^{r+3}\mathbb{Z})$ we have $\sharp\mathrm{Conj}(u^{2^r})=3\cdot 2^2=12$ by Lemma 5.2. Again by Corollary 6.13, we have $\sharp H\cap\mathrm{Conj}(u^2)\leq a(u,2)_{10}+2^9(12-2)=19\cdot 2^{10}$. Hence

$\frac{\sharp H\cap \mathrm{Conj}(u^2)}{\sharp \mathrm{Conj}(u^2)} \leq \frac{19\cdot 2^{10}}{3\cdot 2^{16}} = \frac{19}{3\cdot 2^6}$. By Proposition 6.17, we have $\sharp H \cap \mathrm{Conj}(u^{2^i}) \leq b(u,2)_{11-i} + 2^{8-i}(12-4) = b(u,2)_{11-i} + 2^{11-i}$ for $2 \leq i \leq 7$. Hence

$$\frac{\sharp H \cap \mathrm{Conj}(u^{2^i})}{\sharp \mathrm{Conj}(u^{2^i})} \leq \frac{b(u,2)_{11-i} + 2^{11-i}}{3\cdot 2^{18-2i}}$$

for $2 \leq i \leq 7$. Therefore

$$\begin{aligned}\frac{\sharp I\backslash G/H}{[G:H]} &\leq \frac{2-1}{2}\cdot\frac{23}{3\cdot 2^7} + \frac{2-1}{2^2}\cdot\frac{19}{3\cdot 2^6} + \sum_{i=2}^{7}\frac{2-1}{2^{i+1}}\cdot\frac{b(u,2)_{11-i}+2^{11-i}}{3\cdot 2^{18-2i}} + \frac{1}{2^8}\\ &= \frac{23+19+15+11+7+5+3+2+3}{3\cdot 2^8} = \frac{11}{3\cdot 2^5}.\end{aligned}$$

Consequently, $\delta_H \geq 1 - 3\cdot\frac{11}{3\cdot 2^7} - 4\cdot 0 - 6\cdot\frac{11}{3\cdot 2^5} = \frac{128-11-88}{2^7} > 0$.

$\square$

Proposition 7.11. *Assume $p = 2$. Take a slim subgroup $H \subseteq \mathrm{SL}_2(\mathbb{Z}/2^{10}\mathbb{Z})$. If H/H_1 is contained in F or $H/H_1 = \mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$, then $\delta_H > 0$.*

Proof. In $\mathrm{SL}_2(\mathbb{Z}/2^{10}\mathbb{Z})$ we have $\sharp\mathrm{Conj}(\sigma) = 3\cdot 2^{17}$, $\sharp\mathrm{Conj}(\tau) = 2^{19}$ and $\sharp\mathrm{Conj}(u^{2^i}) = 3\cdot 2^{16-2i}$ for $1 \leq i \leq 6$ by Lemma 5.1, 5.2.

Suppose $H/H_1 \subseteq F$. By Lemma 4.6, we see that in $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ we have $\sharp f_{2,1}^{-1}(F)\cap \mathrm{Conj}(\sigma) = 0$, $\sharp F\cap\mathrm{Conj}(\tau) = 2$, $\sharp f_{2,1}^{-1}(F)\cap\mathrm{Conj}(u) = 0$ and $\sharp f_{2,1}^{-1}(F)\cap\mathrm{Conj}(u^2) = 3$. We have $\sharp(H/H_3)\cap\mathrm{Conj}(\tau) \leq \sharp f_{3,1}^{-1}(F)\cap\mathrm{Conj}(\tau) = 2\cdot 2^4 = 32$ by Lemma 5.3. By Proposition 6.16, we have $\sharp H\cap\mathrm{Conj}(\tau) \leq a(\tau,2)_{10} + 2^8(32-8) = 13\cdot 2^{11}$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} \leq \frac{13\cdot 2^{11}}{2^{19}} = \frac{13}{2^8}$. The same calculation as in the previous lemma shows

$$\frac{\sharp H \cap \mathrm{Conj}(u^{2^i})}{\sharp \mathrm{Conj}(u^{2^i})} \leq \frac{b(u,2)_{10-i} + 2^{10-i}}{3\cdot 2^{16-2i}}$$

for $1 \leq i \leq 6$. Therefore

$$\begin{aligned}\frac{\sharp I\backslash G/H}{[G:H]} &\leq \frac{2-1}{2}\cdot 0 + \sum_{i=1}^{6}\frac{2-1}{2^{i+1}}\cdot\frac{b(u,2)_{10-i}+2^{10-i}}{3\cdot 2^{16-2i}} + \frac{1}{2^7}\\ &= \frac{15+11+7+5+3+2+3}{3\cdot 2^7} = \frac{23}{3\cdot 2^6}.\end{aligned}$$

Consequently, $\delta_H \geq 1 - 3\cdot 0 - 4\cdot\frac{13}{2^8} - 6\cdot\frac{23}{3\cdot 2^6} = \frac{64-13-46}{2^6} > 0$.

Next suppose $H/H_1 = \mathrm{SL}_2(\mathbb{Z}/2\mathbb{Z})$. Since $H/H_2 \subsetneq \mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ by Lemma 2.1, it is conjugate to A_1 by Lemma 4.7. Lemma 4.10 shows that in $\mathrm{SL}_2(\mathbb{Z}/4\mathbb{Z})$ we have $\sharp A_1\cap\mathrm{Conj}(\sigma) = 3$, $\sharp A_1\cap\mathrm{Conj}(\tau) = 2$, $\sharp A_1\cap\mathrm{Conj}(u) = 0$ and $\sharp A_1\cap\mathrm{Conj}(u^2) = 0$. By Proposition 6.14, we have $\sharp H\cap\mathrm{Conj}(\sigma) \leq a(\sigma,2)_{10} + 2^8(3-2) = 73\cdot 2^8$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\sigma)}{\sharp\mathrm{Conj}(\sigma)} \leq \frac{73\cdot 2^8}{3\cdot 2^{17}} = \frac{73}{3\cdot 2^9}$. We have $\sharp(H/H_3)\cap\mathrm{Conj}(\tau) \leq \sharp f_{3,2}^{-1}(A_1)\cap\mathrm{Conj}(\tau) = 2\cdot 2^2 = 8$ by Lemma 5.3. By Proposition 6.16, we have $\sharp H\cap\mathrm{Conj}(\tau) \leq a(\tau,2)_{10} = 5\cdot 2^{12}$. Hence $\frac{\sharp H\cap\mathrm{Conj}(\tau)}{\sharp\mathrm{Conj}(\tau)} \leq \frac{5\cdot 2^{12}}{2^{19}} = \frac{5}{2^7}$. The same calculation as in the case $H/H_1 \subseteq F$ shows

$$\frac{\sharp I\backslash G/H}{[G:H]} \leq \frac{2-1}{2}\cdot 0 + \frac{2-1}{2^2}\cdot 0 + \sum_{i=2}^{6}\frac{2-1}{2^{i+1}}\cdot\frac{b(u,2)_{10-i}+2^{10-i}}{3\cdot 2^{16-2i}} + \frac{1}{2^7} = \frac{31}{3\cdot 2^7}.$$

Consequently, $\delta_H \geq 1 - 3\cdot\frac{73}{3\cdot 2^9} - 4\cdot\frac{5}{2^7} - 6\cdot\frac{31}{3\cdot 2^7} = \frac{512-73-80-248}{2^9} > 0$.

$\square$

Proposition 3.5, Lemma 7.1 and Proposition 7.2 - 7.11 imply Theorem 3.9.

References

[1] P. Deligne, M. Rapoport, *Les schémas de modules de courbes elliptiques.* Modular functions of one variable, II, 143–316. Lecture Notes in Math., Vol. 349, Springer, Berlin, 1973.

[2] B. Edixhoven, *Rational torsion points on elliptic curves over number fields (after Kamienny and Mazur).* Séminaire Bourbaki, Vol. 1993/94. Astérisque No. 227 (1995), Exp. No. 782, 4, 209–227.

[3] G. Faltings, *Finiteness theorems for abelian varieties over number fields.* Translated from the German original [Invent. Math. 73 (1983), no. 3, 349-366; ibid. 75 (1984), no. 2, 381] by Edward Shipz. Arithmetic geometry (Storrs, Conn., 1984), 9-27, Springer, New York (1986).

[4] S. Kamienny, *Torsion points on elliptic curves and q-coefficients of modular forms.* Invent. Math. 109 (1992), no. 2, 221-229.

[5] N. Katz, B. Mazur, *Arithmetic moduli of elliptic curves.* Annals of Mathematics Studies, 108. Princeton University Press, Princeton, NJ, 1985.

[6] D.-S. Kubert, *Universal bounds on the torsion of elliptic curves.* Proc. London Math. Soc. (3) 33 (1976), no. 2, 193-237

[7] J. Manin, *The p-torsion of elliptic curves is uniformly bounded.* Translated from the Russian original [Izv. Akad. Nauk SSSR Ser. Mat. 33 1969 459–465]. Mathematics of the USSR-Izvestija, Vol. 3 (1969), No. 3-4, 433-438.

[8] B. Mazur, *Modular curves and the Eisenstein ideal.* Publ. Math. Inst. Hautes Études Sci. No. 47 (1977), 33-186.

[9] B. Mazur, *Rational points on modular curves.* Modular functions of one variable V, Lecture Notes in Math., Vol. 601, Springer, Berlin (1977), 107-148.

[10] B. Mazur, *Rational isogenies of prime degree (with an appendix by D. Goldfeld).* Invent. Math. 44 (1978), no. 2, 129-162.

[11] L. Merel, *Bornes pour la torsion des courbes elliptiques sur les corps de nombres.* Invent. Math. 124 (1996), no. 1-3, 437–449.

[12] F. Momose, *Rational points on the modular curves $X_{\mathrm{split}}(p)$.* Compositio Math. 52 (1984), no. 1, 115–137.

[13] F. Momose, *Isogenies of prime degree over number fields.* Compositio Math. 97 (1995), no. 3, 329–348.

[14] K. Nakata, *On the 2-adic representation associated to an elliptic curve defined over $\mathbb{Q}$.* (Japanese), Number Theory Symposium in Kinosaki, December 1979, 221-235.

[15] P. Parent, *Bornes effectives pour la torsion des courbes elliptiques sur les corps de nombres.* J. Reine Angew. Math. 506 (1999), 85–116.

[16] P. Parent, *Towards the triviality of $X_0^+(p^r)(\mathbb{Q})$ for $r > 1$.* Compositio Math. 141 (2005), no. 3, 561–572.

[17] J.-P. Serre, *Abelian l-adic representations and elliptic curves.* Lecture at McGill University, New York-Amsterdam, W. A. Benjamin Inc. (1968).

[18] J.-P. Serre, *Propriétés galoisiennes des points d'ordre fini des courbes elliptiques.* Invent. Math. 15 (1972), no. 4, 259-331.

[19] J.-P. Serre, *Représentations l-adiques.* Algebraic number theory (Kyoto Internat. Sympos., Res. Inst. Math. Sci., Univ. Kyoto, Kyoto, 1976), 177–193. Japan Soc. Promotion Sci., Tokyo, 1977.

[20] J.-P. Serre, *Points rationnels des courbes modulaires $X_0(N)$ [d'après B. Mazur].* Séminaire Bourbaki, 30e année (1977/78), Exp. No. 511, 89-100, Lecture Notes in Math., 710, Springer, Berlin, 1979.

[21] G. Shimura, *Introduction to the arithmetic theory of automorphic functions.* Princeton University Press, Princeton, NJ, 1994.

[22] J. Silverman, *The arithmetic of elliptic curves.* Graduate Texts in Mathematics, 106. Springer-Verlag, New York, 1986.

Keisuke Arai, Graduate School of Mathematical Sciences, The University of Tokyo, Tokyo 153-8914, Japan

E-mail address: `araik@ms.u-tokyo.ac.jp`